

A faster algorithm for Minimum Cycle Basis of graphs

(Submitted to Track A)

Telikepalli Kavitha^{1*}, Kurt Mehlhorn^{1*}, Dimitrios Michail^{1*}, Katarzyna Paluch^{2**}

¹ Max-Planck-Institut für Informatik, Saarbrücken, Germany.
{kavitha, mehlhorn, michail}@mpi-sb.mpg.de

² Institute of Computer Science, University of Wrocław, Poland.
abraka@ii.uni.wroc.pl

Abstract. In this paper we consider the problem of computing a minimum cycle basis in a graph G with m edges and n vertices. The edges of G have non-negative weights on them. The previous best result for this problem was an $O(m^\omega n)$ algorithm, where ω is the best exponent of matrix multiplication. It is presently known that $\omega < 2.376$. We obtain an $O(m^2 n + mn^2 \log n)$ algorithm for this problem. When the edge weights are integers, we have an $O(m^2 n)$ algorithm. For unweighted graphs which are reasonably dense ($m \geq n^{1.7}$), our algorithm runs in $O(m^\omega)$ time. For any $\epsilon > 0$, we also design a $1 + \epsilon$ approximation algorithm to compute a cycle basis which is at most $1 + \epsilon$ times the weight of a minimum cycle basis. The running time of this algorithm is $O(\frac{m^\omega}{\epsilon} \log(W/\epsilon))$ for reasonably dense graphs, where W is the largest edge weight.

1 Introduction

1.1 The problem

Let $G = (V, E)$ be a graph. A *cycle* of G is any subgraph in which each vertex has even degree. Associated with each cycle is an *incidence vector* x , indexed on E , where $x_e = 1$ if e is an edge of C , $x_e = 0$ otherwise. The vector space over $GF(2)$ generated by the incidence vectors of cycles is called the *cycle space* of G . It is well-known that when G is connected, this vector space has dimension $N = m - n + 1$, where m is the number of edges of G and n is the number of vertices. A maximal set of linearly independent cycles is called a *cycle basis*.

The edges of G have non-negative weights. The weight of a cycle is the sum of the weights of its edges. The weight of a cycle basis is the sum of the weights of its cycles. We consider the problem of computing a cycle basis of minimum weight in a graph. (We use the abbreviation MCB to refer to a minimum cycle basis.)

1.2 Background

This problem has been extensively studied, both in its general setting and in special classes of graphs. Its importance lies in understanding the cyclic structure of a graph and its use as a preprocessing step in several algorithms. Such algorithms include algorithms for diverse applications like electrical circuit theory [CC73], structural engineering [CHR76], and periodic event scheduling [dP95].

The oldest known references to the minimum cycle basis are Stepanec [Ste64] and Zykov [Zyk69]. Though polynomial time algorithms for this problem were claimed, these algorithms were not

* Partially supported by the Future and Emerging Technologies programme of the EU under contract number IST-1999-14186 (ALCOM-FT).

** Work done while the author was at MPII supported by Marie Curie Doctoral Fellowship.

correct [HS75,Kol80]. The first polynomial time algorithm for the minimum cycle basis problem was given by Horton [Hor87], and had running time $O(m^3n)$.

Horton's approach was to create a set M of mn cycles which he proved was a superset of an MCB and then extract the MCB as the shortest $m-n+1$ linearly independent cycles from M using Gaussian elimination. Golynski and Horton [GH02] observed that the shortest $m-n+1$ linearly independent cycles could be obtained from M in $O(m^\omega n)$ time using fast matrix multiplication algorithms, where ω is the best exponent for matrix multiplication. It is presently known [CW90] that $\omega < 2.376$. The $O(m^\omega n)$ algorithm was the best known algorithm for the MCB problem.

De Pina [dP95] gave an $O(m^3 + mn^2 \log n)$ to compute an MCB in a graph. The approach in [dP95] is different from that of Horton; de Pina's algorithm is similar to the algorithm of Padberg and Rao [PR82] to solve the minimum weighted T -odd cut problem. Our new algorithm to compute an MCB is also based on the same approach.

Related results: For planar graphs, Hartvigsen and Mardon [HM94] showed that an MCB can be computed in $O(n^2 \log n)$ time. In [HM93] Hartvigsen and Mardon study the structure of minimum cycle bases and characterise graphs whose short cycles³ form an MCB.

Closely related to problem of computing an MCB is the problem of finding a minimum *fundamental cycle basis*, i.e., given a connected graph G , find a spanning tree T of G such that the fundamental cycle basis (where each cycle is of the form: some edges of T and one edge from $G \setminus T$) is as small as possible. This problem has been shown to be NP-complete [DPK82]. The minimum cycle basis problem is also NP-complete when negative edge lengths are allowed.

1.3 New Results

In this paper we obtain the following new results.

For graphs with arbitrary non-negative weights on edges, we give an $O(m^2n + mn^2 \log n)$ algorithm to compute an MCB, improving upon the current $O(m^\omega n)$ upper bound. In particular, whenever $m \geq n \log n$, we have an $O(m^2n)$ algorithm. Also, when the edge weights are integers, we have an $O(m^2n)$ algorithm. When the edge weights are small integers (which also includes unweighted graphs) and $m \geq n^{1.7}$, we have an $O(m^\omega)$ algorithm.

We use an all pairs shortest paths (APSP) algorithm as a subroutine in our algorithm. We obtain the better running times for integer edge weights and unweighted graphs by using faster all pairs shortest path algorithms for these cases [Sei95,GM97,Tho99,Tho00]. Similarly, when the graph is sparse, there are faster APSP algorithms, using which, our algorithm can be made faster⁴.

We also look at approximation algorithms for computing a minimum cycle basis in a graph. Given any $\alpha > 1$, we have an α -approximation algorithm by relaxing the shortest paths subroutine to an α stretch paths⁵ subroutine. The running time of our algorithm which computes a cycle basis whose weight is at most twice the weight of an MCB is $\tilde{O}(m^{3/2}n^{3/2}) + O(m^\omega)$ using the result in [CZ01] to compute 2 stretch paths. For graphs with $m \geq n^{1.6}$, this is an $O(m^\omega)$ algorithm. Using the all pairs $1+\epsilon$ stretch paths algorithm [Zwi98], for any $\epsilon > 0$, we have an $\tilde{O}(mn^\omega/\epsilon \log(W/\epsilon)) + O(m^\omega)$ algorithm to compute a cycle basis which is at most $1+\epsilon$ times the weight of an MCB, where W is the largest edge weight in the graph. Assuming that $m \geq n^{1.7}$ and all edge weights are polynomial in n , this is an $O(m^\omega 1/\epsilon \log(1/\epsilon))$ algorithm.

Using the properties of a minimum cycle basis, we also give an $O(m^\omega)$ algorithm to construct a witness of a minimum cycle basis.

³ A cycle C is considered a short cycle if it is the shortest cycle through one of its edges

⁴ Our algorithm cannot be made to run faster than m^ω though.

⁵ An α stretch (s, t) path is a path which is at most α times the length of a shortest (s, t) path.

Organisation of the paper: The rest of this paper is organised as follows. In Section 2 we give a simple algebraic framework (based on De Pina’s algorithm) to compute a minimum cycle basis in a graph and analyse it. In Section 3 we present our improved algorithm. In Section 4 we give an α -approximation algorithm to compute a cycle basis whose weight is $\leq \alpha \cdot$ weight of an MCB. In Section 5 we give an algorithm to obtain a certificate or witness of a minimum cycle basis.

2 A Simple MCB Algorithm

De Pina [dP95] gave a combinatorial algorithm (see Appendix A) to compute a minimum cycle basis in a graph with non-negative weights on its edges. We feel that the intuition behind the algorithm and the idea as to why it works is not clear from the combinatorial version of the algorithm. So, we interpret this algorithm algebraically. From the algebraic version of the algorithm, the scope for improvement is also clear.

2.1 An algebraic interpretation

Let $G = (V, E)$ be an undirected graph with m edges and n vertices and its edges have non-negative weights. We assume that G is connected. Let T be any spanning tree of G . Let e_1, \dots, e_N be the edges of $G \setminus T$ in some arbitrary but fixed order.

A cycle in G can be viewed in terms of its incidence vector and so each cycle is a vector in the space spanned by all the edges (with 0’s and 1’s in its coordinates). Here we will only look these vectors restricted to the coordinates indexed by $\{e_1, \dots, e_N\}$.

In SIMPLE-MCB (see Fig. 1) we compute the cycles of a minimum cycle basis and their *witnesses*. A witness S of a cycle C is a subset of $\{e_1, \dots, e_N\}$ which will prove that C belongs to our minimum cycle basis. We will view these witnesses or subsets in terms of their incidence vectors over $\{e_1, \dots, e_N\}$.

Hence, both cycles and witnesses are vectors in the space $\{0, 1\}^N$. $\langle C, S \rangle$ stands for the standard inner product of the vectors C and S . We say that a vector S is orthogonal to C if $\langle S, C \rangle = 0$. Since we are in the field $GF(2)$, observe that $\langle C, S \rangle = 1$ if and only if C contains an odd number of edges of S . We present in Fig. 1 a succinct description of the algorithm SIMPLE-MCB.

For $i = 1$ to N do the following:

1. Let S_i be any arbitrary non-zero vector in the subspace orthogonal to $\{C_1, C_2, \dots, C_{i-1}\}$. That is, S_i is a non-trivial solution to the set of linear equations:

$$\langle C_k, x \rangle = 0 \text{ for } k = 1 \text{ to } i - 1.$$
 (Initially, S_1 is any arbitrary non-zero vector in the space $\{0, 1\}^N$.)
2. Compute a shortest cycle C_i such that $\langle C_i, S_i \rangle = 1$.

Fig. 1. SIMPLE-MCB: An algebraic framework for computing an MCB

Since each S_i is non-zero, it has to contain at least one edge e from $G \setminus T$. The cycle formed by edges of T and e has intersection of size exactly 1 with S_i . So, there is always at least one cycle with an odd number of edges of S_i .

Note that C_i is independent of C_1, \dots, C_{i-1} because any vector v in the span of $\{C_1, \dots, C_{i-1}\}$ satisfies $\langle v, S_i \rangle = 0$ (since $\langle C_j, S_i \rangle = 0$ for each $1 \leq j \leq i - 1$), whereas $\langle C_i, S_i \rangle = 1$. Hence, it follows immediately that $\{C_1, \dots, C_N\}$ is a basis.

We still have to describe how to compute a shortest cycle C_i such that $\langle C_i, S_i \rangle = 1$ and how to compute a non-zero vector S_i in the subspace orthogonal to $\{C_1, \dots, C_{i-1}\}$. We will do that in Sections 2.2 and 2.3 respectively. We will first prove that $\{C_1, \dots, C_N\}$ computed in SIMPLE-MCB forms an MCB.

Theorem 1. *The set $\{C_1, C_2, \dots, C_N\}$ determined in SIMPLE-MCB is a minimum cycle basis.*

Proof. (from [dP95]) Suppose not. Then there exists an $0 \leq i < N$ such that there is a minimum cycle basis B that contains $\{C_1, \dots, C_i\}$ but there is no minimum cycle basis that contains $\{C_1, \dots, C_i, C_{i+1}\}$. Since the cycles in B form a spanning set, there exist cycles D_1, \dots, D_k in B such that

$$C_{i+1} = D_1 + D_2 + \dots + D_k$$

Since $\langle C_{i+1}, S_{i+1} \rangle = 1$, there exists some D_j in the above sum such that $\langle D_j, S_{i+1} \rangle = 1$. But C_{i+1} is a shortest cycle such that $\langle C_{i+1}, S_{i+1} \rangle = 1$. So the weight of $C_{i+1} \leq$ the weight of D_j .

Let $B' = B \cup \{C_{i+1}\} \setminus \{D_j\}$. It is easy to see that B' is also a basis. And B' has weight at most the weight of B which is a minimum cycle basis. So B' is also a minimum cycle basis. It is easy to show that $\{C_1, C_2, \dots, C_{i+1}\} \subseteq B'$ because by assumption $\{C_1, \dots, C_i\} \subseteq B$ and the cycle D_j that was omitted from B' cannot be equal to any one of C_1, \dots, C_i because $\langle D_j, S_{i+1} \rangle = 1$ whereas $\langle C_j, S_{i+1} \rangle = 0 \forall j \leq i$.

The existence of the basis B' contradicts that there is no minimum cycle basis containing $\{C_1, \dots, C_i, C_{i+1}\}$. Hence, $\{C_1, C_2, \dots, C_N\}$ is indeed a minimum cycle basis. \square

2.2 Computing the cycles

Given S_i , it is easy to compute a shortest cycle C_i such that $\langle C_i, S_i \rangle = 1$ by reducing it to n shortest path computations in an appropriate graph G_i . The following construction is well-known.

G_i is defined from $G = (V, E)$ and $S_i \subseteq E$ in the following manner.

G_i has 2 copies of each vertex $v \in V$. Call them v^+ and v^- .

for every edge $e = (v, u) \in E$ **do**

if $e \notin S_i$ **then**

 Add edges (v^+, u^+) and (v^-, u^-) to the edge set of G_i . {Assign their weights to be the same as e .}


else

 Add edges (v^+, u^-) and (v^-, u^+) to the edge set of G_i . {Assign their weights to be the same as e .}

end if

end for

G_i can be visualised as 2 levels of G (the + level and the - level). Within each level, we have edges of $E \setminus S_i$. Between the levels we have the edges of S_i . Below is an example of G_i when the graph G has 4 vertices $\{1, 2, 3, 4\}$ and 4 edges $\{(1, 2), (1, 4), (2, 4), (3, 4)\}$ and only the edge $(1, 2)$ is in S_i .



Given any v^+ to v^- path p in G_i , we can correspond to it a cycle in G by identifying the vertices and edges in G_i with their corresponding vertices and edges in G . Because we identify both v^+ and v^- with v , the path in G corresponding to p would be a cycle C .

More formally, take the incidence vector of the path p (over the edges of G_i) and obtain an incidence vector over the edges of G by identifying (v^*, u^\dagger) with (v, u) where $*$ and \dagger are $+$ or $-$. Suppose the path p contained more than one copy of some edge(s). (It could have contained both (v^+, u^-) and (v^-, u^+) for some (v, u) .) Then add the number of occurrences of each such edge modulo 2 to obtain an incidence vector over the edges of G .

Let $p = \min_{v \in V}$ shortest (v^+, v^-) path in G_i . We give the proof of the following lemma in Appendix B.

Lemma 1. *p corresponds a shortest cycle C with odd intersection with S_i .*

The computation of the path p can be done by computing n shortest (v^+, v^-) paths (each by Dijkstra's algorithm) in G_i and taking their minimum or by one invocation of an all-pairs-shortest paths algorithm in G_i . This computation takes $O(n(m + n \log n))$ time. In the case when the edge weights are integers or the unweighted case it is better to use faster all-pairs-shortest paths algorithms than run Dijkstra's algorithm n times.

Since we have to compute totally N such cycles C_1, C_2, \dots, C_N , we spend $O(mn(m + n \log n))$ time, since $N = m - n + 1$.

2.3 Computing the subsets

We will now consider the problem of computing the subsets S_i , for $i = 1$ to N . S_i is a non-zero vector in the subspace orthogonal to $\{C_1, \dots, C_i\}$. One way to find a non-zero vector in a subspace is to maintain the whole basis of the subspace. Any vector in that basis will then be a non-zero vector in the subspace.

Initially, $S_j = \{e_j\}$ for all j , $1 \leq j \leq N$. This corresponds to the standard basis of the space $\{0, 1\}^N$. At the beginning of phase i , we have $\{S_i, S_{i+1}, \dots, S_N\}$ which is a basis of the space \mathcal{C}^\perp orthogonal to the space \mathcal{C} spanned by $\{C_1, \dots, C_{i-1}\}$. We use S_i to compute C_i and update $\{S_{i+1}, \dots, S_N\}$ to a basis $\{S'_{i+1}, \dots, S'_N\}$ of the subspace of \mathcal{C}^\perp which is orthogonal to C_i . The update step of phase i is as follows:

For $i + 1 \leq j < N$, let

$$S'_j = \begin{cases} S_j & \text{if } \langle C_i, S_j \rangle = 0 \\ S_j + S_i & \text{if } \langle C_i, S_j \rangle = 1 \end{cases}$$

Lemma 2. *S'_{i+1}, \dots, S'_N form a basis of the subspace orthogonal to C_1, \dots, C_i .*

Proof. We will first show that S'_{i+1}, \dots, S'_N belong to the subspace orthogonal to C_1, \dots, C_i . We know that S_i, S_{i+1}, \dots, S_N are all orthogonal to C_1, \dots, C_{i-1} . Since each S'_j , $i + 1 \leq j \leq N$ is a linear combination of S_j and S_i , it follows that S'_j is orthogonal to C_1, \dots, C_{i-1} . If an S_j is already orthogonal to C_i , then we leave it as it is, i.e., $S'_j = S_j$. Otherwise, we update it as $S'_j = S_j + S_i$. Since both $\langle C_i, S_j \rangle$ and $\langle C_i, S_i \rangle$ are equal to 1, it follows that each S'_j is now orthogonal to C_i also. Hence, S'_{i+1}, \dots, S'_N belong to the subspace orthogonal to C_1, \dots, C_i .

Now we will show that S'_{i+1}, \dots, S'_N are linearly independent. Suppose there is a linear dependence among them. Substitute S'_j 's in terms of S_j 's and S_i in the linear dependence relation. S_i is the only vector that might occur more than once in that relation. So either S_i occurs an even number of times and gets cancelled and we get a linear dependence among S_{i+1}, \dots, S_N or S_i occurs an odd number of times, in which case we get a linear dependence among S_i, S_{i+1}, \dots, S_N . Both the cases contradict the linear independence of S_i, S_{i+1}, \dots, S_N . Hence, S'_{i+1}, \dots, S'_N are linearly independent.

This completes the description of the algorithm SIMPLE-MCB.

Running Time of SIMPLE-MCB: During the update step of phase i , the cost of updating each $S_j, j > i$ is N and hence it is $N(N - i)$ for updating S_{i+1}, \dots, S_N . Since we have N phases, the total cost of maintaining this basis is N^3 , which is like m^3 .

The total running time of the algorithm SIMPLE-MCB, by summing the costs of computing the cycles and witnesses, is $m^3 + mn^2 \log n$. So, independent of which all-pairs-shortest-paths algorithm is used to compute the cycles, the cost of updating the witnesses is the bottleneck.

Note that we needed just one vector from the subspace orthogonal to $\{C_1, \dots, C_i\}$. But the algorithm maintained $N - i$ such vectors: S_{i+1}, \dots, S_N . This was the limiting factor in the running time of the algorithm.

3 Our improvement

The maintenance of the basis of \mathcal{C}^\perp costed us m^2 in each iteration. In order to improve the running time of SIMPLE-MCB, we relax the invariant that S_{i+1}, \dots, S_N form a basis of the subspace orthogonal to C_1, \dots, C_i . Since we need just one vector in this subspace, we can afford to relax this invariant and maintain the correctness of the algorithm.

In SIMPLE-MCB in phase i we updates S_{i+1}, \dots, S_N . Our idea now is to update only those S_j 's where j is close to i and postpone the update of the later S_j 's. During the postponed update, many S_j 's can be updated simultaneously.

We will use a function *extend_cycle_basis* to implement this idea. The function *extend_cycle_basis* works in a recursive manner.

3.1 Computing the Minimum Cycle Basis using *extend_cycle_basis*

We present below the overall algorithm FAST-MCB and the procedure *extend_cycle_basis*. The function *update* will be described in Section 3.2. Recall that the edges e_1, \dots, e_N are the edges of $G \setminus T$, where T is a spanning tree of G .

- Initialize the cycle basis with the empty set and initialize $S_j = \{e_j\}$ for $1 \leq j \leq N$.
- Call the procedure *extend_cycle_basis*($\{\}, \{S_1, \dots, S_N\}, N$).

(A call to *extend_cycle_basis*($\{C_1, \dots, C_i\}, \{S_{i+1}, \dots, S_{i+k}\}, k$) extends the cycle basis by k cycles. C denotes the current partial cycle basis which is $\{C_1, \dots, C_i\}$.)

The procedure *extend_cycle_basis*($C, \{S_{i+1}, \dots, S_{i+k}\}, k$):

- if $k = 1$, compute a shortest cycle that has odd intersection with S_{i+1} .
- if $k > 1$, we use recursion.
 1. we first call *extend_cycle_basis*($C, \{S_{i+1}, \dots, S_{i+\lfloor k/2 \rfloor}\}, \lfloor k/2 \rfloor$) to extend the current cycle basis by $\lfloor k/2 \rfloor$ elements.
 2. we then call *update*($\{S_{i+1}, \dots, S_{i+\lfloor k/2 \rfloor}\}, \{S_{i+\lfloor k/2 \rfloor+1}, \dots, S_{i+k}\}$) to update $\{S_{i+\lfloor k/2 \rfloor+1}, \dots, S_{i+k}\}$. Let $\{T_{i+\lfloor k/2 \rfloor+1}, \dots, T_{i+k}\}$ be the output returned by *update*.
 3. we then call *extend_cycle_basis*($C \cup \{C_{i+1}, \dots, C_{i+\lfloor k/2 \rfloor}\}, \{T_{i+\lfloor k/2 \rfloor+1}, \dots, T_{i+k}\}, \lfloor k/2 \rfloor$) to extend the current cycle basis by $\lfloor k/2 \rfloor$ cycles.

Fig. 2. FAST-MCB: A faster minimum cycle basis algorithm

The procedure *extend_cycle_basis* ($\{C_1, \dots, C_i\}, \{S_{i+1}, \dots, S_{i+k}\}, k$) computes k new elements C_{i+1}, \dots, C_{i+k} of the minimum cycle basis using the subsets S_{i+1}, \dots, S_{i+k} . We maintain the invariant that these subsets are all orthogonal to C_1, \dots, C_i . It first computes $C_{i+1}, \dots, C_{i+\lfloor k/2 \rfloor}$ using $S_{i+1}, \dots, S_{i+\lfloor k/2 \rfloor}$. At this point, the remaining subsets $S_{i+\lfloor k/2 \rfloor+1}, \dots, S_{i+k}$ need not be orthogonal to the new cycles $C_{i+1}, \dots, C_{i+\lfloor k/2 \rfloor}$. So it then updates $S_{i+\lfloor k/2 \rfloor+1}, \dots, S_{i+k}$ so that they are orthogonal to $C_{i+1}, \dots, C_{i+\lfloor k/2 \rfloor}$ and they continue to be orthogonal to $\{C_1, \dots, C_i\}$. Then it computes $C_{i+\lfloor k/2 \rfloor+1}, \dots, C_{i+k}$.

Let us see a small example as to how this works. Suppose $N = 4$. We initialize the subsets $S_i, i = 1, \dots, 4$ and call *extend_cycle_basis*, which then calls itself with only S_1 and S_2 and then only with S_1 and so computes C_1 . Then we update S_2 so that $\langle C_1, S_2 \rangle = 0$ and compute C_2 . Now we simultaneously update S_3 and S_4 which were still at their initial values so that the updated S_3 and S_4 are both orthogonal to C_1 and C_2 . Now we compute C_3 using S_3 and update S_4 and then compute C_4 .

Observe that whenever we compute C_{i+1} , we have the property that S_{i+1} is orthogonal to C_1, \dots, C_i . The difference is the function update which allows us to update many S_j 's simultaneously to be orthogonal to many C_i 's. This simultaneous update enables us to use fast matrix multiplication algorithms which is crucial to the speedup. We describe these steps in detail in the next section.

3.2 The function *update*:

When we call the function *update* ($\{S_{i+1}, \dots, S_{i+\lfloor k/2 \rfloor}\}, \{S_{i+\lfloor k/2 \rfloor+1}, \dots, S_{i+k}\}$), the sets $S_{i+\lfloor k/2 \rfloor+1}, \dots, S_{i+k}$ need not all be orthogonal to the space spanned by $\mathcal{C} \cup \{C_{i+1}, \dots, C_{i+\lfloor k/2 \rfloor}\}$. We know that $S_{i+\lfloor k/2 \rfloor+1}, \dots, S_{i+k}$ are all orthogonal to \mathcal{C} and now we need to ensure that the updated $S_{i+\lfloor k/2 \rfloor+1}, \dots, S_{i+k}$ (call them $T_{i+\lfloor k/2 \rfloor+1}, \dots, T_{i+k}$) are all orthogonal to $\mathcal{C} \cup \{C_{i+1}, \dots, C_{i+\lfloor k/2 \rfloor}\}$.

We will use the sets $T_{i+1}, \dots, T_{i+\lfloor k/2 \rfloor}$ which are the final versions of the subsets $S_{i+1}, \dots, S_{i+\lfloor k/2 \rfloor}$ in order to perform this update, since only when the computation of these sets $T_{i+1}, \dots, T_{i+\lfloor k/2 \rfloor}$ is completed, do we call the function *update* ($i + \lfloor k/2 \rfloor, \lfloor k/2 \rfloor$). We now want to update the sets $S_{i+\lfloor k/2 \rfloor+1}, \dots, S_{i+k}$, i.e., we want to determine $T_{i+\lfloor k/2 \rfloor+1}, \dots, T_{i+k}$ such that for each j in the range for $i + \lfloor k/2 \rfloor + 1 \leq j \leq i + k$

1. T_j is orthogonal to $C_{i+1}, \dots, C_{i+\lfloor k/2 \rfloor}$ and
2. T_j continues to remain orthogonal to C_1, \dots, C_i .

So we define T_j (for each $i + \lfloor k/2 \rfloor + 1 \leq j \leq i + k$) as follows:

$$T_j = S_j + \text{a linear combination of } T_{i+1}, \dots, T_{i+\lfloor k/2 \rfloor}.$$

This makes sure that T_j is orthogonal to the cycles C_1, \dots, C_i because S_j and all of $T_{i+1}, \dots, T_{i+\lfloor k/2 \rfloor}$ are orthogonal to C_1, \dots, C_i . Hence, T_j which is a linear combination of them will also be orthogonal to C_1, \dots, C_i . The coefficients of the linear combination will be chosen such that T_j will be orthogonal to $C_{i+1}, \dots, C_{i+\lfloor k/2 \rfloor}$.

Let

$$T_j = S_j + a_{j1}T_{i+1} + a_{j2}T_{i+2} + \dots + a_{j\lfloor k/2 \rfloor}T_{i+\lfloor k/2 \rfloor}.$$

We will determine the coefficients $a_{j1}, \dots, a_{j\lfloor k/2 \rfloor}$ for all $i + \lfloor k/2 \rfloor + 1 \leq j \leq i + k$ simultaneously. We now look at this problem as a problem in linear algebra.

A Problem in Linear Algebra: Consider the following problem. We are given an invertible $\lfloor k/2 \rfloor \times \lfloor k/2 \rfloor$ matrix X and a $\lfloor k/2 \rfloor \times \lfloor k/2 \rfloor$ matrix Y and we want to find a $\lfloor k/2 \rfloor \times \lfloor k/2 \rfloor$ matrix A such that:

$$(A \ I) \cdot \begin{pmatrix} X \\ Y \end{pmatrix} = 0$$

Here 0 stands for the $k \times \lfloor k/2 \rfloor$ zero-matrix and I stands for the $\lfloor k/2 \rfloor \times \lfloor k/2 \rfloor$ identity matrix.

We need $AX + Y = 0$ or $A = -YX^{-1} = YX^{-1}$ since we are in the field $GF(2)$. We can determine A in time k^ω using fast matrix multiplication and inverse algorithms.

The Implementation of *update*: We want

$$\begin{pmatrix} T_{i+\lfloor k/2 \rfloor+1} \\ \vdots \\ \vdots \\ T_{i+k} \end{pmatrix} = (A \ I) \cdot \begin{pmatrix} T_{i+1} \\ \dots \\ T_{i+\lfloor k/2 \rfloor} \\ S_{i+\lfloor k/2 \rfloor+1} \\ \dots \\ S_{i+k} \end{pmatrix}$$

where A is a $\lfloor k/2 \rfloor \times \lfloor k/2 \rfloor$ matrix whose ℓ th row has the unknowns $a_{j1}, \dots, a_{j\lfloor k/2 \rfloor}$, where $j = i + \lfloor k/2 \rfloor + \ell$. And T_j represents a row with the coefficients of T_j as its row elements.

Let us multiply both sides of this equation with an $N \times \lfloor k/2 \rfloor$ matrix whose columns are the cycles C_{i+1} to $C_{i+\lfloor k/2 \rfloor}$. Then the left hand side is the 0 matrix since each of the vectors $T_{i+\lfloor k/2 \rfloor+1}, \dots, T_{i+k}$ has to be orthogonal to each of $C_{i+1}, \dots, C_{i+\lfloor k/2 \rfloor+1}$. Let

$$\begin{pmatrix} X \\ Y \end{pmatrix} = \begin{pmatrix} T_{i+1} \\ \dots \\ T_{i+\lfloor k/2 \rfloor} \\ S_{i+\lfloor k/2 \rfloor+1} \\ \dots \\ S_{i+k} \end{pmatrix} \cdot (C_{i+1}^T \dots C_{i+\lfloor k/2 \rfloor}^T)$$

where

$$X = \begin{pmatrix} T_{i+1} \\ \dots \\ T_{i+\lfloor k/2 \rfloor} \end{pmatrix} \cdot (C_{i+1}^T \dots C_{i+\lfloor k/2 \rfloor}^T); \quad Y = \begin{pmatrix} S_{i+\lfloor k/2 \rfloor+1} \\ \dots \\ S_{i+k} \end{pmatrix} \cdot (C_{i+1}^T \dots C_{i+\lfloor k/2 \rfloor}^T)$$

Then

$$0 = (A \ I) \cdot \begin{pmatrix} X \\ Y \end{pmatrix}$$

And we have the problem of the preceding paragraph if we show that X is invertible. The matrix

$$X = \begin{pmatrix} \langle T_{i+1}, C_{i+1} \rangle & \dots & \langle T_{i+1}, C_{i+\lfloor k/2 \rfloor} \rangle \\ \langle T_{i+2}, C_{i+1} \rangle & \dots & \langle T_{i+2}, C_{i+\lfloor k/2 \rfloor} \rangle \\ \vdots & \vdots & \vdots \\ \langle T_{i+\lfloor k/2 \rfloor}, C_{i+1} \rangle & \dots & \langle T_{i+\lfloor k/2 \rfloor}, C_{i+\lfloor k/2 \rfloor} \rangle \end{pmatrix} = \begin{pmatrix} 1 & * & \dots & * \\ 0 & 1 & * & * \\ 0 & 0 & 1 & * \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

is an upper diagonal matrix with 1's on the diagonal, since each T_ℓ is orthogonal to all $C_j, j < \ell$. Hence, X is invertible. Thus $A = YX^{-1}$.

Correctness of FAST-MCB: By the implementation of the function *update*, Lemma 3 follows.

Lemma 3. *When $k = 1$, i.e., whenever we call $\text{extend_cycle_basis}(\{C_1, \dots, C_i\}, S_{i+1}, 1)$, S_{i+1} is orthogonal to $\{C_1, \dots, C_i\}$. And S_{i+1} always contains the edge e_{i+1} .*

Hence, just before we compute C_{i+1} , we always have a non-zero vector S_{i+1} orthogonal to $\{C_1, \dots, C_i\}$. The correctness of FAST-MCB follows then from Theorem 1.

3.3 The running time of FAST-MCB

The recurrence of our FAST-MCB algorithm is as follows:

$$T(k) = \begin{cases} \text{cost of computing a shortest odd cycle } C_i \text{ in } S_i & \text{if } k = 1 \\ 2T(k/2) + \text{cost of update} & \text{if } k > 1 \end{cases}$$

Cost of update: The computation of matrices X and Y takes time $mk^{\omega-1}$ using the fast matrix multiplication algorithm. We are multiplying $k/2 \times N$ by $N \times k/2$ matrices. We split the matrices into $2N/k$ square blocks and use fast matrix multiplication to multiply the blocks. Thus multiplication takes time $(2N/k)(k/2)^\omega = O(mk^{\omega-1})$. We can also invert X in $O(k^\omega)$ and we also multiply Y and X^{-1} using fast matrix multiplication in order to get the matrix A .

Using the algorithm described in Section 2.2 to compute a shortest odd cycle C_i in S_i , the recurrence turns into

$$T(k) = \begin{cases} mn + n^2 \log n & \text{if } k = 1 \\ 2T(k/2) + k^{\omega-1}m + k^\omega & \text{if } k > 1 \end{cases}.$$

This solves to $k(mn + n^2 \log n) + k^\omega + k^{\omega-1}m$. Thus $T(m) = m^\omega + m^2n + mn^2 \log n$.

Since $m^\omega < m^2n$, this reduces to $T(m) = m^2n + mn^2 \log n$.

For $m > n \log n$, this is m^2n . For $m \leq n \log n$, this is $mn^2 \log n$.

Theorem 2. *A minimum cycle basis of an undirected weighted graph can be computed in time $O(m^2n + mn^2 \log n)$.*

Our algorithm has a running time of $O(m^\omega + m \cdot n(m + n \log n))$, where the $n(m + n \log n)$ term is the cost to compute all pairs shortest paths. This term can be replaced with a better term when the graph is unweighted or the edge weights are integers or when the graph is sparse.

When the edges of G have integer weights, we can compute all pairs shortest paths in time $O(mn)$ [Tho99,Tho00], that is we can bound $T(1)$ by $O(mn)$. When the graph is unweighted or the edge weights are small integers, we can compute all pairs shortest paths in time $\tilde{O}(n^\omega)$ [Sei95,GM97]. When such graphs are reasonably dense, say $m \geq n^{1.7}$, then the m^ω term dominates the running time of our algorithm.

Theorem 3. *A minimum cycle basis in a graph with integer edge weights can be computed in time $O(m^2n)$. For unweighted graphs which satisfy $m \geq n^{1.7}$, we have an $O(m^\omega)$ algorithm.*

4 An approximation algorithm for Minimum Cycle Basis

The bottleneck in the running time of our minimum cycle basis algorithm is the computation of the shortest cycle C_i such that $\langle C_i, S_i \rangle = 1$. Suppose we relax our constraint that our cycle basis

should have minimum weight and ask for a cycle basis whose weight is at most α times the weight of an MCB. Then can we give a faster algorithm?

We show a positive answer to the above question. For any parameter $\alpha > 1$, we present below an approximation algorithm which computes a cycle basis whose weight is at most α times the weight of a minimum cycle basis. To the best of our knowledge, this is the first time that an approximation algorithm for the MCB problem is being given.

This algorithm is obtained by relaxing $k = 1$ step in the procedure *extend_cycle_basis* of our FAST-MCB algorithm (Fig. 2). In the original algorithm, we computed a shortest cycle C_{i+1} such that $\langle C_{i+1}, S_{i+1} \rangle = 1$. Here, we relax it to compute a cycle D_{i+1} such that $\langle D_{i+1}, S_{i+1} \rangle = 1$ and the weight of D_{i+1} is at most α times the weight of a shortest cycle that has odd intersection with S_{i+1} . The method of updating the subsets S_i would be identical to the way the updation is done in FAST-MCB.

A succinct description of our algorithm is given in Fig. 3.

For $i = 1$ to N do the following:

- Let S_i be any arbitrary non-zero vector in the subspace orthogonal to $\{D_1, D_2, \dots, D_{i-1}\}$ i.e., S_i is a non-trivial solution to the set of equations:
$$\langle D_k, x \rangle = 0 \text{ for } k = 1 \text{ to } i - 1.$$
- Compute a cycle D_i such that $\langle D_i, S_i \rangle = 1$ and the weight of $D_i \leq \alpha \cdot$ the weight of a shortest odd cycle in S_i .

Fig. 3. APPROXIMATE-MCB: An α -approximate MCB

The linear independence of the D_i 's follows from the existence of S_i 's (by using S_i to show that each D_i is linearly independent of $\{D_1, \dots, D_{i-1}\}$). Similarly, note that the subsets $\{S_1, \dots, S_N\}$ are linearly independent since each S_i is independent of $\{S_{i+1}, \dots, S_N\}$ because $\langle S_i, D_i \rangle = 1$ whereas $\langle S_j, D_i \rangle = 0$ for each $j > i$.

4.1 Correctness of APPROXIMATE-MCB

Let $|C|$ denote the weight of cycle C . We need to show that $\sum_{i=1}^N |D_i| \leq \alpha \cdot \text{weight of MCB}$. Let A_i be a shortest odd cycle in S_i . The set $\{A_1, \dots, A_N\}$ need not be linearly independent since the subsets S_i 's were not updated according to the A_i 's. The following lemma was originally shown in [dP95] in order to give an equivalent characterisation of the MCB problem as a maximisation problem. We present a simple proof of the lemma here.

Lemma 4. $\sum_{i=1}^N |A_i| \leq \text{weight of MCB}$.

Proof. We will look at the A_i 's in sorted order i.e., let π be a permutation on $[N]$ such that $|A_{\pi(1)}| \leq |A_{\pi(2)}| \leq \dots \leq |A_{\pi(N)}|$. Let $\{C_1, \dots, C_N\}$ be the cycles of an MCB and let $|C_1| \leq |C_2| \leq \dots \leq |C_N|$. We will show that for each i , $|A_{\pi(i)}| \leq |C_i|$. That will prove the lemma.

We will first show that $\langle C_k, S_{\pi(\ell)} \rangle = 1$ for some k and ℓ with $1 \leq k \leq i \leq \ell \leq N$. Otherwise, the $N - i + 1$ linearly independent vectors $S_{\pi(i)}, S_{\pi(i+1)}, \dots, S_{\pi(N)}$ belong to the subspace orthogonal to C_1, \dots, C_i ; however, this subspace has dimension only $N - i$.

This means that $|A_{\pi(\ell)}| \leq |C_k|$ since $A_{\pi(\ell)}$ is a shortest cycle such that $\langle A_{\pi(\ell)}, S_{\pi(\ell)} \rangle = 1$.

But by the sorted order, $|A_{\pi(i)}| \leq |A_{\pi(\ell)}|$ and $|C_k| \leq |C_i|$. This implies that $|A_{\pi(i)}| \leq |C_i|$. \square

It is obvious from the algorithm that for each i , $|D_i| \leq \alpha \cdot |A_i|$. Hence, it follows from the above lemma that $\sum_{i=1}^N |D_i| \leq \alpha \cdot \text{weight of MCB}$. Thus Theorem 4 follows.

Theorem 4. *The cycles $\{D_1, \dots, D_N\}$ computed in Fig. 3 have weight at most α times the weight of a minimum cycle basis.*

4.2 The running time of APPROXIMATE-MCB

Since all the steps of APPROXIMATE-MCB, except the step corresponding to computing a shortest cycle, are identical to FAST-MCB, we have the following recurrence for APPROXIMATE-MCB:

$$T(k) = \begin{cases} \text{cost of computing an } \alpha \text{ stretch odd cycle } D_i \text{ in } S_i & \text{if } k = 1 \\ 2T(k/2) + \text{cost of update} & \text{if } k > 1 \end{cases}$$

So the running time of APPROXIMATE-MCB depends on which parameter α is used in the algorithm. We will compute an α stretch odd cycle D_i by using the same method as in Section 2.2. But instead of a shortest (v^+, v^-) path in G_i , here we would compute an α stretch (v^+, v^-) path. It is easy to see that the minimum of such paths would correspond to an α stretch cycle in G that has odd intersection with S_i .

When $\alpha = 2$, we use the result in [CZ01] to compute 2 stretch paths which would result in 2 stretch cycles. Then APPROX-MCB runs in time $\tilde{O}(m^{3/2}n^{3/2}) + O(m^\omega)$. For graphs with $m \geq n^{1.6}$, this is an $O(m^\omega)$ algorithm.

For $1 + \epsilon$ approximation, we use the all pairs $1 + \epsilon$ stretch paths algorithm [Zwi98]. Then we have an $\tilde{O}(mn^\omega/\epsilon \log(W/\epsilon)) + O(m^\omega)$ algorithm to compute a cycle basis which is at most $1 + \epsilon$ times the weight of an MCB, where W is the largest edge weight in the graph. Assuming that $m \geq n^{1.7}$ and all edge weights are polynomial in n , this is an $O(m^\omega 1/\epsilon \log(1/\epsilon))$ algorithm.

5 Computing a Certificate of Optimality

Given a set of cycles $\mathcal{C} = \{C_1, \dots, C_N\}$ we would like to construct a certificate to verify the claim that \mathcal{C} forms an MCB. A certificate is an “easy to verify” witness of the optimality of our answer.

For example, the sets S_i , $1 \leq i \leq N$ in the algorithm from which we calculate the cycles $\mathcal{C} = \{C_1, \dots, C_N\}$ of the minimum cycle basis, are a certificate of the optimality of \mathcal{C} . The running time of the verification algorithm would have smaller constants than FAST-MCB and it would be conceptually much simpler. This motivates the following question: given a set of cycles $\{C_1, \dots, C_N\}$, compute its certificate.

The algorithm below computes witnesses S_1, \dots, S_N given C_1, \dots, C_N .

1. Compute a spanning tree T . Let $\{e_1, \dots, e_N\}$ be the edges of $G \setminus T$.
2. Form the 0-1 $N \times N$ matrix $\mathcal{C} = (C_1^T \dots C_N^T)$, where the i -th column of \mathcal{C} is the incidence vector of C_i over $\{e_1, \dots, e_N\}$.
3. Compute \mathcal{C}^{-1} . The rows of \mathcal{C}^{-1} are our witnesses or certificate.

If the inversion algorithm returns an error, it means that \mathcal{C} is singular, then $\{C_1, \dots, C_N\}$ are linearly dependent, hence they cannot form a cycle basis.

The rows of \mathcal{C}^{-1} form our witnesses S_1, S_2, \dots, S_N . The property that we want from S_1, \dots, S_N is that for each i , $\langle C_i, S_i \rangle = 1$. Since $\mathcal{C}^{-1}\mathcal{C}$ is the identity matrix, this property is obeyed by the rows of \mathcal{C}^{-1} .

Suppose each C_i is a shortest cycle such that $\langle C_i, S_i \rangle = 1$. Then by Lemma 4, this means that $\sum_{i=1}^N |C_i| \leq \text{weight of an MCB}$. Since $\{C_1, \dots, C_N\}$ are linearly independent (by the existence of \mathcal{C}^{-1}), it means that $\{C_1, \dots, C_N\}$ forms a minimum cycle basis.

On the other hand, if for some i , C_i is not a shortest cycle such that $\langle C_i, S_i \rangle = 1$, then by replacing C_i with a shortest cycle that has odd intersection with S_i (as in the proof of Theorem 1), we get a cycle basis with smaller weight.

Hence, the cycles $\{C_1, \dots, C_N\}$ form an MCB if and only if each C_i is a shortest odd cycle in S_i . Since the inverse of an $N \times N$ matrix can be computed in $O(N^\omega)$ time, we have the following theorem.

Theorem 5. *Given a set of cycles $\mathcal{C} = \{C_1, \dots, C_N\}$ we can construct a certificate $\{S_1, \dots, S_N\}$ in $O(m^\omega)$ time.*

References

- [CC73] L. O. Chua and L. Chen. On optimally sparse cycle and coboundary basis for a linear graph. In *IEEE Trans. Circuit Theory*, volume CT-20, pages 495–503, 1973.
- [CHR76] A. C. Cassell, J. C. Henderson, and K. Ramachandran. Cycle bases of minimal measure for the structural analysis of skeletal structures by the flexibility method. In *Proc. Royal Society of London Series A*, volume 350, pages 61–70, 1976.
- [CW90] D. Coppersmith and S. Winograd. Matrix multiplications via arithmetic progressions. *Journal of Symb. Comput.*, 9:251–280, 1990.
- [CZ01] E. Cohen and U. Zwick. All-pairs small-stretch paths. *Journal of Algorithms*, 38:335–353, 2001.
- [dP95] J.C. de Pina. *Applications of Shortest Path Methods*. PhD thesis, University of Amsterdam, Netherlands, 1995.
- [DPK82] N. Deo, G. M. Prabhu, and M. S. Krishnamoorthy. Algorithms for generating fundamental cycles in a graph. *ACM Trans. Math. Software*, 8:26–42, 1982.
- [GH02] Alexander Golynski and Joseph D. Horton. A polynomial time algorithm to find the minimum cycle basis of a regular matroid. In *8th Scandinavian Workshop on Algorithm Theory*, 2002.
- [GM97] Z. Galil and O. Margalit. All pairs shortest paths for graphs with small integer length edges. *Journal of Computing Systems and Sciences*, 54:243–254, 1997.
- [HM93] David Hartvigsen and Russell Mardon. When do short cycles generate the cycle space? *Journal of Combinatorial Theory, Series B*, 57:88–99, 1993.
- [HM94] David Hartvigsen and Russell Mardon. The all-pairs min cut problem and the minimum cycle basis problem on planar graphs. *Journal of Discrete Mathematics*, 7(3):403–418, 1994.
- [Hor87] J. D. Horton. A polynomial-time algorithm to find a shortest cycle basis of a graph. *SIAM Journal of Computing*, 16:359–366, 1987.
- [HS75] E. Hubicka and M. M. Syslo. Minimal bases of cycles of a graph. In M. Fiedler, editor, *Recent Advances in Graph Theory*, pages 283–293, 1975.
- [Kol80] E. Kolasinska. On a minimum cycle basis of a graph. *Zastos. Mat.*, 16:631–639, 1980.
- [PR82] Padberg and Rao. Odd minimum cut-sets and b-matchings. *Mathematics of Operations Research*, 7:67–80, 1982.
- [Sei95] R. Seidel. On the all-pairs-shortest-path problem in unweighted undirected graphs. *Journal of Computing Systems and Sciences*, 51:400–403, 1995.
- [Ste64] G. F. Stepanec. Basis systems of vector cycles with extremal properties in graphs. *Uspekhi Mat. Nauk*, 19:171–175, 1964.
- [Tho99] M. Thorup. Undirected single-source shortest paths with positive integer weights in linear time. *Journal of the ACM*, 46:362–394, 1999.
- [Tho00] M. Thorup. Floats, integers, and single source shortest paths. *Journal of Algorithms*, 35:189–201, 2000.
- [Zwi98] U. Zwick. All pairs shortest paths in weighted directed graphs - exact and approximate algorithms. In *Proc. of the 39th Annual IEEE FOCS*, pages 310–319, 1998.
- [Zyk69] A. A. Zykov. *Theory of Finite Graphs*. Nauka, Novosibirsk, 1969.

Appendix A

De Pina's combinatorial algorithm

De Pina gave the following combinatorial algorithm (Fig.4) to compute a minimum cycle basis in a graph G with non-negative weights on its edges.

Let T be a fixed spanning tree. Let e_1, \dots, e_N be the edges of $G \setminus T$ in some arbitrary but fixed order.

Initialize $S_{1,i} = \{e_i\}$ ($i = 1, \dots, N$).

for $k = 1, \dots, N$ **do**

 Find a shortest cycle C_k with an odd number of edges in $S_{k,k}$

 Define for $i = k + 1, \dots, N$:

$$S_{k+1,i} = \begin{cases} S_{k,i} & \text{if } C_k \text{ has an even number of edges in } S_{k,i} \\ S_{k,i} \Delta S_{k,k} & \text{if } C_k \text{ has an odd number of edges in } S_{k,i} \end{cases}$$

end for

The algorithm returns $\{C_1, \dots, C_N\}$.

Fig. 4. De Pina's combinatorial algorithm for computing a MCB

Since each set $S_{k,k}$ always contains the edge e_k , the cycle formed by edges of T and e_k has intersection exactly 1 with $S_{k,k}$. Hence, the set of cycles with an odd number of edges in $S_{k,k}$ is non-empty.

Appendix B

Proof of Lemma 1

Proof. The path p has to contain an odd number of edges of S_i . This is because only edges of S_i provide a change of sign and p goes from a $+$ vertex to a $-$ vertex. We might have deleted some edges of S_i while forming C since those edges occurred with a multiplicity of 2. But this means that we always delete an even number of edges from S_i . Hence C has an odd number of edges of S_i present in it. Also, the length of $C \leq$ the length of p since edges have non-negative lengths.

We should now prove that C is a shortest such cycle. Let C' be any other cycle in G with an odd number of edges of S_i in it. If C' is not a simple cycle, then C' is a union of simple cycles (with disjoint edges) and at least one of those simple cycles C_0 should have an odd number of edges of S_i present in it. And the weight of $C_0 \leq$ the weight of C' .

Let u be a vertex in C_0 . We will identify C_0 with a path in G_i by traversing C_0 starting at the vertex u and identifying it with u^+ . If we traverse an edge e of S_i , then we identify the vertices incident on e with opposite signs. And if we traverse an edge outside S_i , then we identify the vertices incident on e with the same sign. Since C_0 is a cycle, we come back to the vertex u . Also, C_0 has an odd number of edges of S_i present in it. So the sign of the final vertex is of the opposite sign to the sign of the starting vertex. Hence, C_0 translates to a u^+ to u^- path p' in G_i . And the length of $p' =$ the length of C_0 .

But p was the minimum weight path among all shortest (v^+, v^-) paths in G_i for all $v \in V$. Hence, the length of $p \leq$ the length of p' . So we finally we get that the length of $C \leq$ the length of $p \leq$ the length of $p' \leq$ the length of C' . This proves that C is a shortest cycle with odd intersection with S_i . \square