

An improved algorithm for online rectangle filling

Rob van Stee*

November 16, 2010

Abstract

We consider the problem of scheduling resource allocation where a change in allocation results in a changeover penalty of one time slot. We assume that we are sending packets over a wireless channel of uncertain and varying capacity. In each time slot, a bandwidth of at most the current capacity can be allocated, but changing the capacity has a cost, which is modeled as an empty time slot. Only the current bandwidth and the bandwidth of the immediately following slot are known. We give an online algorithm with competitive ratio 1.753 for this problem, improving over the previous upper bound of 1.848. The main new idea of our algorithm is that it attempts to avoid cases where a single time slot with a nonzero allocation is immediately followed by an empty time slot. Additionally, we improve the lower bound for this problem to 1.6959. Our results significantly narrow the gap between the best known upper and lower bound.

Keywords: resource management and awareness, wireless networks, online algorithms

1 Introduction

In wireless networks, channel conditions can change frequently, which affects the bit error rate and therefore the channel transmission capacity [5]. We consider the problem of setting data transmission rates over such a channel in order to maximize the throughput. Naturally, at any time the transmission rate cannot be higher than the current transmission capacity, but there is also typically a nonzero cost involved in changing the transmission rate, because the transmitter and receiver will have to coordinate and reset to a new transmission rate. We model this cost as the loss of a single time slot. That is, whenever we want to change the transmission rate (or if we are forced to change it, because the current capacity is below the rate that we set earlier), we will have one time slot in which nothing can be transmitted.

Formally, we are given an online sequence of nonnegative real numbers $h(1), h(2), \dots$, which represent the maximum transmission capacities of the wireless channel at each time step, and we need to determine the transmission rate $u(i)$ at each time step. Our goal is to maximize $\sum_i u(i)$, and due to the changeover cost we have for any i that $u(i) = u(i+1)$, $u(i) = 0$, or $u(i+1) = 0$. It can be seen that if only the current bandwidth is known, no competitive online algorithm exists [2]. We therefore focus on the case where some information about future bandwidth is given; in particular, for our results we assume that we have a lookahead of a single time slot.

The name rectangle filling comes from a geometrical interpretation of the problem, where each time slot i is represented by a rectangle of unit width and height $h(i)$ (also called a *column*). An algorithm needs to

*Research supported by German Research Foundation (DFG). Department of Computer Science, University of Karlsruhe, Germany. vanstee@ira.uka.de

decide how much of each rectangle to fill, i.e., what the transmission rate $\in [0, h(i)]$ should be. Any feasible solution for this problem is a set of rectangles (of varying width) where the transmission rate is constant; all these rectangles are separated by one or more zero columns.

Probabilistic analysis for this problem was given by Tsibonis et al. [6] and Borst [3]. Arora and Choi were the first ones to study this problem from a worst-case perspective [1]. They gave a dynamic program for the offline version with a running time of $O(n^3)$, and a 4-competitive online algorithm called Wait Dominate Hold (using a lookahead of 1). Arora et al. [2] soon afterwards showed that this algorithm is actually $8/3$ -competitive, and gave a lower bound of $8/5$. For the version of the problem with k -lookahead, they gave an online algorithm with competitive ratio 2 for any k , and a lower bound of $(k + 2)/(k + 1)$.

Wang et al. [7] presented a faster offline algorithm with a running time of $O(n^2)$, and new lower and upper bounds of 1.6358 and 1.848 respectively. In the following year, the same authors [8] considered the version with k -lookahead. They gave a deterministic algorithm with a competitive ratio of $1 + 2/(k - 1)$, as well as a randomized algorithm with competitive ratio $1 + 1/(k + 1)$. They also gave a randomized lower bound of $1 + 1/(\sqrt{k + 2} + \sqrt{k + 1})^2$, which is more than $1 + 1/(4k + 8)$ and tends to $1 + 1/(4k + 4)$ for large k . The randomized lower bound was recently improved to $(k + 2) \ln \frac{k+2}{k+1} > 1 + 1/(2k + 3)$ by Epstein and Levin [4].

Generally, despite the seeming simplicity of the model, the gap between the upper and lower bounds has so far remained relatively large, particularly for the most basic version with a lookahead of 1, and it appears to be very hard to give tight bounds for this problem. See also the conclusions.

Our results We give an improved algorithm which achieves a competitive ratio of less than 1.753. The main new idea of our algorithm is to try and limit the amount of changes in the used bandwidth. Therefore, as soon as a nonzero transmission rate is allocated, we temporarily relax the condition for changing the rate, and allow a *single* time slot with relatively high capacity to appear without resetting the transmission rate. The idea behind this is that as long as there is only a single slot with high capacity, we do not lose too much compared to the optimal solution, because the optimal solution needs to allocate zero columns before and after the slot in order to be able to use the full capacity of the slot. That is, it is not worth the trouble of paying a penalty to serve a slot completely, especially if we have very recently paid another penalty to start transmitting.

The analysis uses the natural block partitioning from Wang et al. [7] as a starting point, but is significantly more involved. Apart from their partitioning rules, we will need a number of additional assignment rules to deal with the columns that are left unassigned in their scheme. These assignment rules depend on the optimal solution, and sometimes assign part of the optimal profit to preceding blocks and part to following blocks in order to allow us to analyze these blocks independently.

Moreover, in some cases we analyze the competitive ratio by splitting the instance into two parts, replacing one column by a sequence of columns, and showing that our algorithm gives exactly the same allocation as before to all columns where it does not allocate 0, and the optimal profit is split according to another set of rules between the first part and the second part of the input.

Finally, we improve the lower bound from 1.6358 to 1.6959. The construction is similar to the one from Wang et al. [7], but we use an extra threat in every step of the input.

2 Algorithm MoreFilling

We first give an informal description of our algorithm. MoreFilling creates blocks between zero columns. The nonzero part of a block starts by trying to guess a good allocation for the first two nonzero columns.

The first column arrives at time $t = 1$. We take $u(0) = 0$.

1. If $u(t-1) > h(t)$, set $u(t) = 0$.
2. If $u(t-1) = 0$, set $u(t)$ depending on $h(t+1)/h(t)$, as follows:

$$\frac{h(t+1)/h(t)}{u(t)} \quad \left| \quad \begin{array}{cccc} [0, \beta) & [\beta, 1) & [1, \gamma) & [\gamma, \infty) \\ h(t) & h(t+1) & h(t) & 0 \end{array} \right.$$

3. If $u(t-1) > 0$, let $t_1 = 1 + \max\{t \geq 0 \mid u(t) = 0\}$, and set $H = \min(h(t_1), h(t_1 + 1))$.
 If $t_1 = t - 1$ and $h(t)/h(t-1) \in [1, \delta]$, set $q_3 := \gamma$, else $q_3 := \delta$.
 If $h(t+1) \geq q_3 H$, set $u(t) = 0$, else $u(t) = u(t-1)$.

Figure 1: The algorithm MoreFilling

If the first height is much smaller than the second one (by a factor of at least $\gamma > 2$, then the nonzero part of the block is simply postponed till later. Otherwise, if the first height is much larger (by a factor of at least $\frac{1}{\beta} > 1.4$, then the first column is allocated its full height, and the block will contain a single nonzero column. If the two heights are relatively close, the minimum between the two heights is allocated.

The main idea of our algorithm is to try and avoid having zero columns if the nonzero part of a block has just started. Hence there is one case where the first nonzero column is fully used, where we allow the column height to grow by a factor of $\gamma > 2$ for just one step, while still keeping other (later) column heights bounded by a smaller factor. This ensures that there is only a zero column if the new column height is significantly larger, or if the nonzero column is allocated its full height.

The third column in a block is almost always limited to a maximum allowed height of $\delta < 1.6$. The only exception is if the second column has height between 1 and δ times the first column, i.e., the block does not contain a column of height (almost) γ yet. We never let the heights grow by a factor of γ if the block already contains at least two nonzero columns.

We now formally define our algorithm. Define the following values.

$$\begin{array}{llll} \mathcal{R} = & & = 1.75214 & \beta = \mathcal{R}/(2\mathcal{R} - 1) = 0.69966 \\ \varepsilon = & (2\mathcal{R} - 3)\beta & = 0.35282 & \gamma = \mathcal{R}/(4 - 2\mathcal{R} + \varepsilon) = 2.06489 \\ \delta = & (2\mathcal{R}\beta - \varepsilon - 1)/\beta & = 1.57074 & \eta = 1 - \mathcal{R}/\gamma + \varepsilon = 0.50414 \end{array}$$

We have $(5 - \mathcal{R}/\gamma + \varepsilon)/(1 + \delta) = \mathcal{R}$. Our algorithm is defined in Figure 1.

Theorem 1 *The competitive ratio of MoreFilling is at most $\mathcal{R} = 1.75214$.*

It should be pointed out that δ could be set to any value between roughly 1.55 and $5/3$, and MoreFilling would still be 1.753-competitive. The other variables, β, ε and γ are decisive.

3 Analysis

We begin our analysis by introducing the block partitioning and giving some properties of the optimal solution in Section 3.1. We use this to analyze the most basic type of block in Theorem 2. We denote an optimal allocation by $a(\cdot)$ and a particular type of difficult column by r . We then consider what happens if

$a(r) = 0$ or $a(r - 1) = 0$ in Section 3.2 (apart from one difficult case), and the case $a(r) = a(r - 1)$ is treated in Section 3.4.

3.1 Block partitioning

We classify the columns with zero allocation (also called zero columns) by the rule in which they are set to 0: a column is of type i if it is set to 0 by Rule i . The zero columns partition the input into blocks in a natural way. Note that type 2 columns only occur after other zero columns. We follow the partitioning scheme from Wang et al. [7], which we describe next.

A type 1 column ends a block, and is a part of that block. If column i is a type 3 column, one block ends at column $i - 1$ and the next starts at column $i + 1$. (In this case, we will decide later what to do with column i .) This partitioning scheme ignores the type 2 columns (which only occur after other zero columns). Each block hence consists of zero or more type 2 columns, followed by one or more nonzero columns and possibly one final type 1 column. For each block B , the number of *nonzero* columns in B , also called the *length* of B , is denoted by $|B|$.

Let us consider the possible optimal profit on a block. We normalize the height of the columns in this block such that the height of the first column is exactly 1.

Definition 1 We define the **BASEHEIGHT** of a block with at least two nonzero columns as the minimum height among its first two columns.

The **BASEHEIGHT** is abbreviated by H in the algorithm.

Definition 2 A block is called *long* if it has at least one (nonzero) column after its first fully-used column, else it is called *short*.

Observation 1 Consider a block of which the first nonzero column is column i . We have $h(i) = 1$. If $h(i) > h(i + 1)$, then $h(i + 1) \geq \beta h(i)$, or $u(i + 1) = 0$. If $u(i + 1) > 0$ and the last column k of this block is of type 1, we have $h(k) < \beta$.

If $h(i) \leq h(i + 1)$, then $h(i + 1) < \gamma$. Moreover, almost always we have $h(j) < \delta$ for all $j > i + 1$ such that j and i belong to the same block. The only exception to this is if $h(i + 1) \leq \delta$, in which case we have $h(i + 2) < \gamma$ if $i + 2$ is part of the same block. In addition, if the last column k of the block of i is of type 1, we have $h(k) < \alpha$.

We now consider the type 2 zero columns at the start of the block.

Lemma 1 (Wang et al.) Given a sequence of columns S , if each column's height is at least γ times the height of the previous one (where $\gamma \geq 2$), then the value of the optimal solution is at most $\frac{\gamma^2}{\gamma^2 - 1}h$ where h is the height of the last column in S . This value is achieved by using every other column completely starting from the right.

In order to efficiently deal with all the cases, we will in fact use the following estimates, which are all higher than the bound from Lemma 1. Consider a column i of height h which is preceded by type 2 columns. We make two distinctions: one based on whether $a(i) > h/\gamma$ or not (if $a(i) > h/\gamma$, then $a(i - 1) = 0$), and one based on whether the block containing column i is long or short (Definition 2). The bounds used are as follows.

	$a(i) > h/\gamma$	$a(i) \leq h/\gamma$
Short block	$\varepsilon \approx 0.353$	$\varepsilon\gamma \approx 0.729$
Long block	$\eta \approx 0.505$	$\varepsilon\gamma \approx 0.729$




Figure 2: This figure shows the maximum possible heights of nonzero columns in a block (plus a final type 1 column). There are three cases. Let the first nonzero column of the block be t , then we have the cases $\delta h(t) < h(t+1) \leq \gamma h(t)$ (left), $h(t) \leq h(t+1) \leq \delta h(t)$ (middle), and $\beta h(t) \leq h(t+1) < h(t)$ (right). If $h(t+1) < \beta h(t)$, the block contains only a single nonzero column; if $h(t+1) > \gamma h(t)$, no block starts at time t . The third column may have height γ only if the heights are ascending and the second column has height at most δ . Shown is the case where the sixth column is of type 1; if there are additional nonzero columns instead, their maximum height is the same as that of the fifth column.

Naturally the profit on type 2 columns does not really depend on the type of the following block, but this assumption simplifies the analysis later.

Observation 2 *If the input contains a sequence of columns i, \dots, j such that $h(k) \geq \gamma h(k-1)$ for $k = i+1, \dots, j$, then MoreFilling earns 0 on columns $i, \dots, j-1$.*

To determine the maximum optimal profit on a block, we need to consider how the type 2 blocks at the start of such a block are serviced. Depending on the exact heights of the nonzero columns, it may not be optimal to service them as described in Lemma 1. However, Observation 2 allows us to make the following assumption.

Assumption 1 *If columns i, \dots, j form a sequence of type 2 columns followed by a nonzero column j , then $h(k) = \gamma h(k-1)$ for $k = i+1, \dots, j+1$.*

Here we simply round up column heights, which can only improve the total optimal value. Regarding MoreFilling, if the previous block ended with a type 1 column, its behavior is unaffected. If it ended with a type 3 column, then the column following that was too high for the block to continue, and it is now not less high than before.

Fix an optimal solution and denote its allocation to column i by $a(i)$.

Lemma 2 *For each column i , we have $a(i) = 0$ or $a(i) > h(i)/3$.*

Proof Suppose the optimal allocation for columns $i-1, i, i+1$ is $x, a(i), z$ (if one of these columns does not exist, assume that its height is zero), then we have $x = 0$ or $x = a(i)$ and we also have $z = 0$ or $z = a(i)$, so if $a(i) < h(i)/3$ we can replace $x, a(i), z$ by $0, h(i), 0$, which is always feasible. \square

We have the following lemma which will later simplify the analysis.

Lemma 3 *Let S be a sequence of type 2 columns. Let the height of the final type 2 column be h . Let $V(S)$ be the optimal profit on S after adding $\eta = 1 - \mathcal{R}/\gamma + \varepsilon$ times the allocation to the first column. If the allocation to the first column is nonzero, then $V(S) \leq \varepsilon \gamma^2 h$.*

Proof By Lemma 2 and Assumption 1, for any column $i \geq 1$ in S , only three allocations then need to be considered: $0, h(i)/\gamma$, and $h(i)$.

Let x be the number of columns. If x is odd, the optimal allocation to S is given by Lemma 1, and this satisfies the constraint that the first column have nonzero allocation. For $x = 1$, the optimal profit on S is h . To this we add ηh to get $V(S)$, and we have $1 + \eta = \varepsilon\gamma^2$. For any odd $x \geq 3$ we have

$$V(S) = h + \frac{h}{\gamma^2} + \cdots + \frac{h}{\gamma^{x-1}} \left(2 - \frac{\mathcal{R}}{\gamma} + \varepsilon\right)$$

while for $x + 2$ this value becomes

$$h + \cdots + \frac{h}{\gamma^{x-1}} + \frac{h}{\gamma^{x+1}} \left(2 - \frac{\mathcal{R}}{\gamma} + \varepsilon\right)$$

which is less since $\gamma^2 \left(2 - \frac{\mathcal{R}}{\gamma} + \varepsilon\right) > \gamma^2 + 2 - \frac{\mathcal{R}}{\gamma} + \varepsilon$. Therefore, $V(S) \leq \varepsilon\gamma^2 h$ for all odd x .

For $x = 2$, given that the first column (of height $\frac{h}{\gamma}$) must have nonzero allocation, it is best to assign the same value to the second column, and we find $V(S) \leq \frac{1}{\gamma^2} \left(3 - \frac{\mathcal{R}}{\gamma} + \varepsilon\right) < \varepsilon\gamma^2 h$ by inspection.

Given an instance with x type 2 columns (x is even), consider the instance with $x + 2$ type 2 columns. Let the height of column x be h' . If it was optimal to use column x fully, then it is now optimal to use column $x + 2$ (of height $\gamma^2 h'$) fully as well. This adds $\gamma^2 h'$ to the total profit. The profit on the first x columns cannot be improved (by induction on even x), and there is no other assignment to column x that increases the profit on the last two columns to a value above $\gamma^2 h'$. Thus, we do not need to consider other allocations to column x in this case. Analogously to the case of odd x , we can now show that $V(S)$ decreases for increasing x . \square

Theorem 2 *MoreFilling is \mathcal{R} -competitive on a block which is **not** immediately followed by a type 3 column.*

Proof Let FIRST be the number of the first nonzero column in block b , and normalize $h(\text{FIRST}) = 1$. Then, the profit of our algorithm on this block is $|b|$ or $\beta|b|$, depending on whether $h(\text{FIRST}) \leq h(\text{FIRST} + 1)$. By the text below Lemma 1, the optimal profit on the type 2 columns in b is at most $\varepsilon\gamma h(\text{FIRST})$ (the height of the last type 2 column is at most $h(\text{FIRST})/\gamma$, and if the last type 2 column is not used, the optimal profit on the other type 2 columns is at most $\varepsilon h(\text{FIRST})$, or at most $\eta h(\text{FIRST})$ if the block is long). For the calculations, we make the worst-case assumption that there is a type 1 column at the end of this block, and it has height 1 (it actually must have smaller height). The exception to this is a block of length 1; in that case, the type 1 column must have height at most $\beta h(\text{FIRST})$ by our algorithm.

Note that all nonzero columns in the block have height at least 1. To derive upper bounds for the optimal profit on a block, we may assume that each of these columns has the maximum height as bounded by Observation 1 (Figure 2). The only allocations that we need to consider to find the optimal profit are values that are the height of at least one column. In particular, for column FIRST , by Lemma 2 only the following allocations need to be considered: $h(\text{FIRST})$, $\frac{1}{\gamma} h(\text{FIRST})$, and possibly $\beta h(\text{FIRST})$.

This gives us the results shown in Figure 3. The figure shows the optimal profit for each case and compares it to the profit of MoreFilling. Regarding a block of length 1 for instance, we find that it is optimal to set $a(\text{FIRST}) = a(\text{FIRST} + 1) = h(\text{FIRST} + 1) \leq \beta h(\text{FIRST})$, $a(\text{FIRST} - 1) = 0$, and then it is possible to earn at most $\varepsilon h(\text{FIRST}) < 0.353h(\text{FIRST})$ on the type 2 columns up to column $\text{FIRST} - 2$.

For a block that contains three or more nonzero columns, note that adding a column of height δ (resp. $\delta\beta$ in case $h(\text{FIRST} + 1) < h(\text{FIRST})$) adds at most δ (resp. $\delta\beta$) to the optimal profit and exactly 1 (resp. β) to the profit of MoreFilling. Since $\delta < \mathcal{R}$, this does not increase the competitive ratio above \mathcal{R} . \square

To complete our analysis, we now need to eliminate the type 3 columns.

Let column r be a type 3 zero column. These columns are the most difficult to handle for the following reason. For type 1 columns, MoreFilling already decided in the previous time step (or earlier) to allocate zero to the type 1 column, because its height is too small compared to some previous height. Similarly, type




Figure 3: Possible profiles of blocks that are followed by type 1 columns. Left are blocks which start with a fully-used column, right are blocks where the second column is fully-used. The triangles indicate preceding type 2 columns (that have geometrically increasing heights), of value at most $\varepsilon < 0.353$, or $\eta < 0.505$ for long blocks. The dashed blocks indicate blocks created by MoreFilling. Bold lines indicate the optimal solution for a block.

2 columns are allocated zero because their height is small compared to the immediately following height. For these cases, it is clear how to group the columns into blocks as described above (i.e., how to compensate for the missed profit on the zero column), and we can analyze these cases in a straightforward way as shown in Theorem 2.

In contrast, type 3 columns are allocated zero “at the last minute”, and this decision does *not* depend on its own height. This is in particular troublesome in the case where column $r - 1$ was allocated $h(r) < h(r - 1)$. In normal cases, column r would be allocated $h(r)$ to compensate for the fact that less than $h(r - 1)$ was earned on column $r - 1$. But now, column r is allocated zero, and in many cases we will have to consider the block following column r to complete the analysis.

Again, we normalize the heights of the columns such that the height of the first nonzero column of the preceding block BEFORE is 1. Then $h(r) \leq \gamma$, and if the first column of BEFORE is not fully used or the second column has height at least δ , we have $h(r) \leq \delta H$, where H is the base height of block BEFORE (else column $r - 1$ would have been zeroed). We also have that

$$h(r + 1) > \delta H. \quad (1)$$

Denote the block following column r by AFTER. Suppose a counterexample I exists, that shows a competitive ratio above \mathcal{R} . We next introduce a sequence of assignment rules to deal with type 3 columns.

3.2 The cases $a(r) = 0$ and $a(r - 1) = 0$

Assignment rule 1 *If $h(r + 1) \geq \gamma h(r)$, we assign column r as a type 2 zero column to AFTER. Additionally, assign $\varepsilon h(r)$ of the optimal profit on BEFORE to AFTER; if $a(r) = 0$, assign $\varepsilon \gamma h(r)$.*

As long as we have $|\text{BEFORE}| \geq 2$ or $h(r) \geq h(r - 1)$, this works because the allocation of MoreFilling to column $r - 1$ was not adversely affected by column r . That is, $u(r - 1)$ was not set to a value below $h(r - 1)$ because it was followed by a column of height $h(r)$ (but possibly because it was preceded by other columns with nonzero allocation), and thus block BEFORE can be analyzed independently of column r , i.e., as if the input ended at column $r - 1$. (We can just ignore the fact that some of the optimal profit on BEFORE was assigned to AFTER in this case, and apply Theorem 2 on BEFORE.)

Moreover, the profit of $\varepsilon h(r)$ (or $\varepsilon \gamma h(r)$ if $a(r) = 0$) is a lower bound for what we assume the optimal solution could earn on AFTER from additional type 2 columns *before* column r (i.e. of heights $h(r)/\gamma, h(r)/\gamma^2, \dots$ (from right to left)). Since these extra type 2 columns are not in fact there, we can reassign this profit from BEFORE to AFTER.

Suppose $|\text{BEFORE}| = 1$ and $h(r) < h(r-1)$. We show that also in this case, MoreFilling is \mathcal{R} -competitive on BEFORE after applying this assignment rule. Normalize such that $h(r-1) = \text{MF}(\text{BEFORE}) = 1$. Then $\text{MF}(\text{BEFORE}) = h(r) \geq \beta$. We need to show that $\text{OPT}(\text{BEFORE}) < \mathcal{R}\beta = 1.22591$.

If $a(r-1) = 0$, we have (after the reassignment) $\text{OPT}(\text{BEFORE}) \leq \varepsilon\gamma h(r-1) - \varepsilon h(r) < 1 - \varepsilon\beta < 1$.

If $a(r) = 0$, we have

$$\text{OPT}(\text{BEFORE}) \leq h(r-1) + \varepsilon h(r-1) - \varepsilon\gamma h(r) \leq 1 + \varepsilon - \varepsilon\gamma\beta < 1.$$

Else, $a(r-1) = a(r) > 0$. In this case, if $a(r-2) = 0$ we have

$$\text{OPT}(\text{BEFORE}) \leq 1 + \varepsilon - \varepsilon h(r) \leq 1 + \varepsilon - \varepsilon\beta < 1.11$$

and otherwise $\text{OPT}(\text{BEFORE}) \leq \frac{1}{\gamma} + \varepsilon\gamma - \varepsilon h(r) < \frac{1+\varepsilon}{\gamma} < 1$.

Assignment rule 2 *If after applying Assignment rule 1 there is a type 3 column r in I for which $a(r) = 0$, then if $|\text{BEFORE}| \geq 2$ or $h(r) \geq h(r-1)$, we set $h(r) = 0$ and assign column r as a type 1 column to BEFORE. If $|\text{BEFORE}| = 1$ and $h(r) < h(r-1)$, assign column r to BEFORE and assign $\delta\varepsilon h(r)$ of the optimal profit on BEFORE to AFTER.*

We explain why this rule works. Consider a column r for which $a(r) = 0$. First of all, note that the type (or the height) of a zero column does not affect the behavior of MoreFilling on subsequent columns.

If $|\text{BEFORE}| \geq 2$ or $h(r) \geq h(r-1)$, our modification does not change the profit of MoreFilling on the modified block BEFORE, and the optimal value for I does not decrease since $a(r) = 0$. Thus MoreFilling is competitive on BEFORE (including column r , seen as a type 1 column) by Theorem 2.

Suppose $|\text{BEFORE}| = 1$ and $h(r) < h(r-1) = 1$. By rule 3 of our algorithm, we have $h(r+1) > \delta h(r)$. On hypothetical type 2 columns preceding column $r+1$, OPT could hence earn at least $\varepsilon h(r+1) > \delta\varepsilon h(r)$, and we assign $\delta\varepsilon h(r)$ of profit to AFTER. Now $\text{MF}(\text{BEFORE}) = h(r) \geq \beta$, and using $a(r) = 0$ we get

$$\text{OPT}(\text{BEFORE}) \leq (1 + \varepsilon)h(r-1) - \delta\varepsilon h(r) = 1 + \varepsilon - \delta\varepsilon h(r) \leq 1 + \varepsilon - \delta\varepsilon\beta < 1 < \mathcal{R}\beta.$$

After applying these two assignment rules, we have

$$a(r) > 0 \quad \text{for any type 3 column } r \text{ in } I$$

3.3 The case $|\text{BEFORE}| = 1$ and $a(r-1) = a(r) = a(r+1) > 0$

We treat this special case separately because we will encounter a very similar case later.

Case 1: $h(r) < h(r-1)$ Consider column $r+1$. If it is of type 2, then we must have $a(r+2) = 0$ by Lemma 2, since $h(r+2) > \gamma h(r+1) > \gamma\delta h(r) \geq \gamma\delta a(r) > 3a(r)$. But the total optimal profit on columns r and $r+1$ is $2a(r)$, whereas by Lemma 1 the total possible profit on type 2 columns preceding column $r+2$ is $\frac{\gamma^2}{\gamma^2-1}h(r+1) > 2.05h(r+1) \geq 2.05a(r)$. So we can assign these columns to AFTER.

If column $r+1$ is nonzero, then the profit of $2a(r)$ is less than $(1 + \varepsilon)h(r+1)$ because that is at least $(1 + \varepsilon)\delta a(r) = 2.12a(r)$. Hence we now find a smaller optimal profit in all cases where it was previously optimal to assign the first column of the block its full height, or where it was assigned zero. For the remaining cases, we have $a(r) \leq h(r) \leq \beta$. On BEFORE, MF earns at least $h(r)$, so we can assign $\mathcal{R}h(r)$ of profit there, in particular $(\mathcal{R} - 1)h(r) - \varepsilon$ from column r , and are then left with at most $(2 - \mathcal{R})h(r) - \varepsilon$ for AFTER. In all these remaining cases, in the optimal solutions which we found before, the first and second column of the block were both used, giving a profit of $2\beta + \varepsilon$ on those columns and the preceding type 2 columns, which in the current situation is at least $(2\beta + \varepsilon)\delta h(r) = 2.752h(r)$ using that $h(r+1) > \delta h(r)$. Now, we find a profit of at most $(2 - \mathcal{R})h(r) - \varepsilon + 2h(r) = 2.248h(r) - \varepsilon$ there which is less.




Figure 4: The missing case. We show the first four subcases of $h(r) \geq h(r-1)$. The first two diagrams represent $h(r+2) \geq \delta$, the next diagram the case $r+2$ is of type 1, and the last that $r+2$ is nonzero.

Case 2: $h(r) \geq h(r-1)$ If $h(r) < \min(h(r+1), \gamma)$, we may increase $h(r)$ to this value without affecting the behavior of MoreFilling or decreasing the optimal profit. Then $h(r) > \delta$.

If now we no longer have $a(r-1) = a(r) = a(r+1) > 0$ (i.e., this is no longer optimal), we are done. Else, if $a(r+2) = 0$, we have a contradiction, since it is better to set $a(r-1) = 0$ and earn more than $2\delta > 3$ on the columns $r, r+1$.

If $h(r+2) \geq \delta$, we find analogously $a(r+3) = a(r+2) = a(r+1)$. If column $r+2$ is of type 1, MoreFilling earns at least $\delta/\beta + 1$ on columns $r-1$ and $r+1$ for a competitive ratio of less than 1.25 on these two blocks. Else, if $h(r+3) \geq \delta$ as well, we have a contradiction in any event, since we could earn $4\delta > 6$ by setting $a(r-1) = 0$ and $a(r+4) = 0$, which is more than we get now on columns $r-1, \dots, r+4$. Therefore $h(r+3) < \delta < h(r+1)$, and column $r+3$ is of type 1, regardless of whether column $r+1$ is nonzero or of type 2.

If column $r+1$ is of type 2, then $\text{MF}(\text{BEFORE} \cup \text{AFTER}) > 1 + \delta\gamma > 4$ (Figure 4.1). If column $r+1$ is nonzero, we have $\text{OPT}(\text{BEFORE} \cup \text{AFTER}) = 5 + \varepsilon$ and $\text{MF}(\text{BEFORE} \cup \text{AFTER}) \geq 1 + 2\delta > 4$ (Figure 4.2). The ratio is less than 1.5 in both cases.

We are left with the case $h(r+2) < \delta$. If $h(r+2) < \delta\beta$, column $r+2$ is of type 1, and the competitive ratio on $\text{BEFORE} \cup \text{AFTER}$ is at most

$$\frac{\varepsilon + 4}{1 + \delta} = 1.694 \quad (\text{Figure 4.3}).$$

Else, if column $r+2$ is nonzero, we get a ratio of at most

$$\frac{\varepsilon + 5}{1 + 2\beta\delta} = 1.674 \quad (\text{Figure 4.4})$$

(less if $|\text{AFTER}| > 2$). Finally, if column $r+2$ is of type 3, we have $h(r+3) > \delta h(r+2) > \delta^2\beta = 1.726$ whereas $a(r+2) = a(r-1) \leq 1$. On hypothetical type 2 columns preceding column $r+3$, OPT could earn $\varepsilon\gamma h(r+3) > 1$. Therefore we can assign $a(r+2)$ completely to the block following AFTER , and have a competitive ratio of

$$\frac{\varepsilon + 3}{1 + \beta\delta} = 1.598$$

on the blocks BEFORE and AFTER .

3.4 The case $a(r) = a(r-1) > 0$

For the remaining cases, we now introduce a set of rules that split the hypothetical counterexample I in two parts, and show that MoreFilling is \mathcal{R} -competitive on both parts.

Case	Condition	$a(r)$	Optimal profit on S	Profit assigned to I_1
1	$a(r-1) = a(r), f = 2, \text{BEFORE} = 2$	β	$\varepsilon h(r+1) > \varepsilon \delta \beta$	$\beta(1 - \varepsilon \delta)$
2	$a(r-1) = a(r), f = 1, \text{BEFORE} = 1,$ $a(r+1) = 0$	1	$\varepsilon \gamma h(r+1) > \varepsilon \gamma \delta$	0
3	$a(r-1) = a(r), \text{other cases}$	δH	$\varepsilon h(r+1) > \varepsilon \delta H$	$\delta(1 - \varepsilon)H$
4	$a(r+1) = a(r-1) = 0, \text{BEFORE} = 1$	γ	$\varepsilon \gamma h(r+1) > \varepsilon \gamma \delta$	$\gamma(1 - \varepsilon \delta)$
5	$a(r-1) = 0, \text{BEFORE} > 1$	γ	$\varepsilon h(r+1) > \varepsilon \delta$	$\gamma - \varepsilon \delta$

Table 1: Replacement rules. The third column in the table contains an upper bound on $a(r)$ for each case. The fourth column contains the optimal profit on S , where we have assumed that the schedule on the remaining columns (all columns but column r) is unchanged. In particular, if $a(r+1) = a(r) > h(r)/\gamma$, the final column in S cannot be used in such a schedule, but if $a(r+1) = 0$, it can be used, thus multiplying its value for the optimal solution by γ . Finally, the final column is an upper bound for the difference between the two preceding columns.




Figure 5: Possible profiles of blocks for the case $a(r-1) = a(r)$. For an explanation of the symbols, see Figure 3. The rectangles marked S are not assigned to the block, but instead to the following block. See Replacement rules. In the leftmost diagram, the case where $a(r+1) = 0$ is shown (Replacement rule 2). The case of a block of one column that is not fully allocated can be handled by Assignment rule 2.

Replacement rules Split I into two independent parts, I_1 and I_2 . Part I_1 ends with column $r-1$. Replace column r by an infinite sequence S of columns of heights $\min(\gamma a(r), h(r+1))/\gamma^i$ ($i = 1, 2, \dots$). Part I_2 consists of sequence S followed by columns $r+1$ and beyond. All columns of S are of type 2. Note that $h(r+1) > \delta \cdot \text{BASEHEIGHT}$. Finally, add some amount to the optimal profit on I_1 as described in the following table. Let $f = 2$ if the second nonzero column of BEFORE has height less than the first one, else $f = 1$. The first column of BEFORE has height 1, and we abbreviate its BASEHEIGHT by H . See Table 1.

Case 1: $a(r-1) = a(r), f = 2, |\text{BEFORE}| = 2$ See Figure 5 (third diagram).

Case 2: $a(r-1) = a(r), f = 1, a(r+1) = 0, |\text{BEFORE}| = 1$ In this case, the optimal profit on S is at least $\varepsilon \gamma \delta a(r) > a(r)$, so nothing is left to assign to I_1 . See Figure 5 (first diagram). Note that the alternative subcase $a(r+1) > 0$ is already handled by Assignment Rule 3.

Case 3: $a(r-1) = a(r), \text{other cases}$ If BEFORE has no fully-used column, then $\text{OPT}(\text{BEFORE}) \leq \mathcal{R}$ by Assignment Rule 1. Else, we have $h(r) \leq \delta$, or column $r-1$ would be a zero column by Rule 3 of our algorithm. Since $a(r) \leq h(r) \leq \delta$, we then only need to assign $\delta(1 - \varepsilon)$ of profit to I_1 , even if $a(r+1) > 0$. See Figure 5 for the optimal schedules in the various cases. As usual, it can be seen that adding additional columns of height δ does not increase the competitive ratio.




Figure 6: Possible profiles of blocks for the case $a(r-1) = 0$. Black triangles are type 2 columns of total value at most $\varepsilon\gamma < 0.729$. In the third diagram, the shown bold allocation is not in fact optimal, because it is better to set $a(r-1) = a(r) = \beta$. Thus, this case is not relevant. Finally, longer blocks do not give a ratio above \mathcal{R} .

Case 4: $a(r+1) = a(r-1) = 0, |\text{BEFORE}| = 1$ Note that the profit assigned to I_1 is less than 1, and there is now no type 1 column at the end of BEFORE. We use that $a(r-1) = 0$. If $|\text{BEFORE}| = 1$ and $u(r-1) = h(r-1)$, then since $a(r-1) = 0$, the optimal profit is now at most $\varepsilon\gamma + \gamma(1 - \varepsilon\delta) = 1.649$, and MoreFilling earns 1. See Figure 6 (top left).

Case 5: $a(r-1) = 0, |\text{BEFORE}| > 1$ In this case we can only be sure of a profit of $\varepsilon\delta$ that can be assigned to I_2 . However, this is sufficient in these cases. See Figure 6 (other diagrams).

This only leaves one case open, which is handled in the following section.

3.5 The final case

Due to our calculations so far, we only still need to deal with the following case:

$$|\text{BEFORE}| = 1, u(r-1) = h(r-1), a(r-1) = 0 \text{ and } a(r+1) = a(r) > 0. \quad (2)$$

In this unresolved case, the first question we have is how much profit must be assigned forward (relative to the online allocation to AFTER). Let x be the number of type 2 columns immediately following column r , and let r_2 be the zero column immediately following $|\text{AFTER}|$.

Lemma 4 *The profit that is assigned forward from column r in case (2) is at most $(1 - \frac{\mathcal{R}}{\gamma} + \varepsilon)a(r) = \eta a(r)$. Normalizing such that the first nonzero column of AFTER has height 1, this is at most η/γ^x .*

Proof We have $a(r) \leq \min(h(r), h(r+1))$. We assign $(\mathcal{R}/\gamma - \varepsilon)a(r) \leq (\mathcal{R}/\gamma - \varepsilon)h(r) \leq (\mathcal{R} - \varepsilon\gamma)h(r-1)$ to BEFORE (to add to the $\varepsilon\gamma h(r-1)$ from the type 2 columns) and are left with $(1 - \frac{\mathcal{R}}{\gamma} + \varepsilon)a(r) \leq (1 - \frac{\mathcal{R}}{\gamma} + \varepsilon)h(r+1)$. The lemma follows because the sequence of type 2 columns has (at least) geometrically increasing heights. In particular, if $x = 0$, we use that $a(r) \leq h(r+1)$, and $h(r+1) = 1$ in this case. \square

Using Lemma 3, we have that if $x \geq 1$, we get exactly the value $\varepsilon\gamma$ for the type 2 columns plus this forward assignment from column r , which is the value that we have been calculating with in case the last type 2 column had nonzero allocation in the optimal solution. Thus, we get exactly the same analysis as before, with the difference (for $x = 1$) that OPT is required to use the type 2 column by assumption. Clearly, if we remove this requirement (and generally use the bound $\varepsilon > \eta/\gamma$ in case OPT does not use the final type 2 column), we can only get higher bounds for the competitive ratio. Hence, our previous analysis holds unless $x = 0$, i.e., there are no type 2 columns before AFTER. (Of course, if for AFTER we encounter the open case that we are discussing in this section, we just repeat; eventually we reach the end of the sequence or find a case which we can prove.) In the case $x = 0$, the optimal solution must use the first nonzero column of AFTER (as well as the preceding column). We summarize this discussion in the following lemma.

Lemma 5 *In the remaining open case, column $r + 1$ is the first nonzero column of block AFTER, and $a(r + 1) = a(r)$.*

Lemma 6 *MoreFilling is competitive on AFTER even after taking the forward assignment from column r into account, unless $|\text{AFTER}| = 1$ and $a(r - 1) = a(r) = a(r + 1)$.*

Proof If AFTER is a long block, we already calculated with an optimal profit of at least η on the type 2 columns in all cases and are done. Else if AFTER is followed by a type 3 column r_2 , we see in Figures 5 and 6 that MoreFilling is still \mathcal{R} -competitive if the profit on preceding columns is η instead of ε (or $\varepsilon\gamma$) apart from in the excluded case. If AFTER is not followed by a type 3 column, then if $|\text{AFTER}| = 1$ we get a ratio of at most $(2\beta + \eta\beta) = \mathcal{R}$, using that $a(r + 1) = a(r) \leq \beta$ and Lemma 4. If $|\text{AFTER}| = 2$ and $f = 2$, then if $a(r + 1) = a(r + 2)$ as in Figure 3, we find a ratio of at most $(3a(r) + \eta a(r))/(2a(r)) = \mathcal{R}$ since now $a(r) \leq u(r + 2)$. Else, the ratio is at most $(\eta + 1 + \beta)/(2\beta) < 1.6$. \square

We are left with the case $|\text{AFTER}| = 1$, so $r_2 = r + 2$ and $a(r - 1) = \dots = a(r + 2)$ since type 3 columns have nonzero optimal allocation. Since r_2 is of type 3,

$$h(r_2 + 1) > \delta h(r_2 - 1) = \delta h(r + 1) \geq \delta a(r). \quad (3)$$

If $a(r_2 + 1) = 0$, we apply our Replacement rule 2, i.e., column r_2 is assigned completely to the block following AFTER. This works because on (hypothetical) type 2 columns preceding column $r_2 + 1$, OPT could earn $\varepsilon\gamma h(r_2 + 1) > h(r_2 + 1)/\delta > a(r_2)$. We then have $\text{OPT}(\text{AFTER}) \leq 1 + (1 - \frac{\mathcal{R}}{\gamma} + \varepsilon) < 1.51$.

Else, we are again in the case discussed in Section 3.3, since $a(r_2 - 1) = a(r_2) = a(r_2 + 1)$. The only difference is that the profit from previous columns is not ε but η . We can therefore follow that analysis completely and adjust the calculations where necessary. The highest ratio that we found there is $(4 + \varepsilon)/(1 + \delta)$, and this now becomes $(4 + \eta)/(1 + \delta) = \mathcal{R}$.

4 Lower bound

We let $\varepsilon > 0$ be some very small value. Denote the online algorithm by ALG and assume that it has a competitive ratio of at most $(1 - \varepsilon)1.69595$. In our lower bound construction, the adversary uses the following strategy. See Figure 7. We have $p < 1$ and $q > 2$. Each rectangle represents a state. The states consist of three components: the current online allocation, the current column, and the next column. Hence, if we move from one state to another, the second column in the starting state is identical to the first column in the destination state. However, we sometimes (conceptually) rescale column heights, for instance when we move from state B to state A .

Additionally, the number $(1 - \delta)x$ in the ellipse indicates that one final column of height $(1 - \delta)x$ arrives, where x is the (nonzero) allocation that ALG just used. Hence, this column has no value for ALG (it can only use the column by forfeiting a profit of x on the current column).

The input The input starts in state A . The default state changes are given by the normal arrows. As ALG is processing the input, moving between states A and B , the adversary keeps track of the total online profit so far, as well as of the total optimal profit. When going from state B to A , the adversary may elect to change the value of q to $q' = 2.036442$ and p to $p' = 1/q'$ (for future columns); we will specify when this happens. In both state A and B , nonstandard state changes are possible, which are indicated by dotted lines in Figure 7. The adversary forces such a state change as soon as this gives the desired lower bound. Note that in all of the cases where a dotted line is followed, the input consists of at most two more columns, and it is straightforward to calculate the resulting competitive ratio, based on the stored optimal and online profits.




Figure 7: Lower bound strategy. The variable x always indicates the allocation chosen by ALG. At the state change marked with (*), we *may* change q to $q' = 2.03644$ and p to $p' = 1/q'$.

For instance, if ALG allocates a certain value $x > 0$ in state A or B , and the adversary sees that this is too low to maintain the competitive ratio (because of earlier choices), the input sequence stops immediately. In state B , this may also happen when the algorithm allocates 0; hence, there is no condition next to the outgoing dotted line from B . In state A , instead of stopping immediately, one final column of height $(1 - \delta)x$ may arrive, so the adversary has two options here (besides the default state change to B) if ALG allocates a nonzero value.

What we need to show is that no matter what ALG does, at some point the adversary can choose a nonstandard state change and prove the desired competitive ratio: ALG cannot stay in states A and B forever.

Output of the adversary We first describe how the adversary handles the input as long as ALG stays in A and B . This is very simple: in state A , it allocates 0 if ALG allocates $x > 0$, and alternates between the full column height and 0 otherwise (ending with 0 when ALG allocates a nonzero value). In state B , it always allocates the full column height, except for one special case when the input is about to end. In that case, the allocation in state B will be the same as on the columns following it.

Claim 1 *This output is a feasible solution.*

Proof We only need to show that the adversary does not allocate two different nonzero values to adjacent columns. Given the above rules, this could only happen at a state change. In a move from A to B , the adversary always assigns 0. In a move from B to A (say at column i), the adversary allocates the full column height, and *may* also do this on column $i + 1$, depending on whether there is an odd or an even number of consecutive state A columns following. But columns i and $i + 1$ have the same height here. \square

In our construction, we will set $q = 2.14447$ and

$$p = \frac{3q^2 - 5 + \sqrt{9q^4 - 14q^2 + 9}}{8q^2 - 8} = 0.709039. \quad (4)$$

Claim 2 *For each visit of ALG to state A and every $\varepsilon > 0$, the number of consecutive steps in which ALG can allocate 0 before the adversary will send it to C is bounded.*

Proof Consider the i th visit of ALG to state A . Let the online and optimal profit *before* this visit be ALG_i and OPT_i , respectively ($\text{ALG}_1 = \text{OPT}_1 = 0$). We want to calculate the competitive ratio if the adversary sends ALG to C when ALG assigns 0 after staying in A for N time steps, N even (state change I in Figure

7). We rescale all the column heights such that the columns in state C have heights exactly 1 and $p < 1$. During the steps that ALG spends in state A on its i th visit, the column heights increase exponentially. Hence, for each $i \geq 1$ and each real $\delta > 0$, there exists an N_δ such that $\text{ALG}_i \leq \delta$ for $N \geq N_\delta$.

On the columns $1, \dots, n_1 - 1$, the adversary earns at least $\text{OPT}_i + \sum_{j=1}^{N/2} q^{-2j}$. This tends to $\text{OPT}_i + 1/(q^2 - 1)$ for large N . Hence, for every $i \geq 1$ and each real $\delta > 0$, there exists an N'_δ such that the adversary earns at least $(1 - \delta)/(q^2 - 1)$ for $N \geq N'_\delta$.

Let $N \geq \max(N_\delta, N'_\delta)$. In state C , ALG must choose an allocation x . If $x > p$ or $x = 0$, the input stops. The total online profit is at most $1 + \text{ALG}_i \leq 1 + \delta$, and the optimal profit is at least $2p + (1 - \delta)/(q^2 - 1)$. If $0 < x \leq p$, the next and final column has height $(1 - \delta)x$. The online profit is at most $2x + \text{ALG}_i \leq 2x + \delta$, whereas the optimal profit is at least $3(1 - \delta)x + (1 - \delta)/(q^2 - 1)$. As a function of x , the competitive ratio in this case is minimized by taking $x = p$. For $\delta \rightarrow 0$, the competitive ratios tend to

$$\mathcal{R}_1 = \frac{3p + 1/(q^2 - 1)}{2p} \quad \text{and} \quad \mathcal{R}_2 = 2p + 1/(q^2 - 1).$$

These ratios are equal if we set p as in (4), and are then both $\mathcal{R} = 1.69595$. Both ratios tend to this value as $\delta \rightarrow 0$. Hence, for any given $\varepsilon > 0$ and $i \geq 1$, we can find $\delta > 0$ and $N \geq \max(N_\delta, N'_\delta)$ such that the implied competitive ratio is at least $(1 - \varepsilon)\mathcal{R}$ if ALG spends at least N steps in state A on its i th visit and the adversary then sends it to C . \square

Claim 3 *For each visit of ALG to state B and every $\varepsilon > 0$, the number of consecutive steps in which ALG can allocate x before the adversary will send it to C (state change IV in Figure 7) is bounded.*

Proof In each step in state B except for the last one, given the above strategy of the adversary, the ratio of the adversary's profit to the online profit is at least $q > 2$. Thus, after sufficiently many consecutive columns in state B (depending on the profit of ALG and the adversary before the current visit to B), the overall competitive ratio gets arbitrarily close to $q > 2 > 1.69595$. \square

Phases Given the previous two claims, we know that ALG must move back and forth between states A and B . We partition the input into phases. A new phase starts at the start of the input and whenever ALG arrives in state A from state B . We say that a column i is of state A (B) if ALG is in state A (B) when it allocates the value for column i . Hence a phase always ends with a state B column in which ALG allocates 0, unless the input stops in state B . In each phase i , ALG allocates exactly one nonzero value (namely when it moves from state A to state B), possibly to some consecutive columns.

Definition 3 *Let $0 < x_i \leq 1$ be the fraction of the height of the first column to which ALG allocates a nonzero value in phase i .*

Claim 4 *In each phase but the last, the adversary earns at least q times as much as ALG. This profit ratio is monotonically nondecreasing as a function of the number of state A columns in the phase.*

Proof For a given phase j , scale the column heights in this phase so that the last column (say column i) in state A has height 1. Let the number of columns in state B be $b \geq 1$. Then ALG earns $bx_j \leq b$ whereas the adversary earns at least bq , namely on columns $i + 1, \dots, i + b$. The end of this phase looks as follows.

Allocations that cause state changes are underlined.

Input	...	q^{-3}	q^{-2}	q^{-1}	1	q	...	q	q
State	A	A	A	A	A	B	...	B	B
ALG	...	0	0	0	x_j	x_j	...	x_j	0
OPT	...	q^{-3}	0	q^{-1}	0	q	...	q	q
									b columns

This proves that the profit ratio is at least q and grows as a function of the number of state A columns. \square

Corollary 1 *W.l.o.g., ALG never stays in state B for more than one column.*

Proof On all state B columns but the last, the adversary earns at least $q > 2$ times as much as ALG as can be seen in the table above. If ALG has a competitive ratio of at most $\mathcal{R} < 2$, then it certainly also has this if it always leaves state B immediately. Thus ignoring extra state B columns can only help ALG; note that removing or ignoring such columns does not affect the column heights of future columns. \square

Let $\text{ALG}(j)$ ($\text{OPT}(j)$) be the total profit of ALG (OPT) after j phases, scaled such that the last state A column in phase j has height 1.

Claim 5 *As $j \rightarrow \infty$, $\text{OPT}(j) \rightarrow q^3/(q^2 - 1) = 2.74$ or more.*

Proof The claimed amount is reached if phase j is arbitrarily long as can be seen in the table in the proof of Claim 4. If phase j is shorter, there are some state B columns, which means that some earlier columns have higher heights. This can only increase the optimal profit. \square

Lemma 7 *For all $j \geq 1$, if ALG is \mathcal{R} -competitive, we have $x_j > q/(2\mathcal{R}) > 0.6$.*

Proof We consider the competitive ratio if the input stops on a nonzero allocation x_j in state A (state change III in Figure 7). The adversary earns q on the final column of the input, whereas ALG earns at most another x_j there. By Claim 4, the competitive ratio is at least

$$\frac{q + q \cdot \text{ALG}(j-1)}{2x_j + \text{ALG}(j-1)} \geq \mathcal{R} \text{ for } x_j \leq \frac{q}{2\mathcal{R}}.$$

(For $j = 1$, we have $\text{ALG}(j-1) = 0$.) \square

We now first deal with the case where after an arbitrary prefix, ALG never spends two consecutive steps in the same state, thus from some point onwards each phase is of the form AB . We then get an input and output of the following form (using the adversary strategy defined above). Here we normalize such that the first column height in phase n is 1.

Input	q^{1-n}	q^{2-n}	q^{2-n}	q^{3-n}	...	q^{-1}	1	1	q
State	A	B	A	B	...	A	B	A	B
Phase	1		2		...	$n-1$		n	
ALG	$x_1 q^{1-n}$	0	$x_2 q^{2-n}$	0	...	$x_{n-1} q^{-1}$	0	x_n	0
OPT	0	q^{2-n}	0	q^{3-n}	...	0	1	0	q

The following observation is immediate from the table and $x_i \in (0, 1]$.

Observation 3 *If ALG allocates a nonzero amount (not necessarily the full column height!) whenever it is in state A , and zero whenever it is in state B , it earns at most $\sum_{i=0}^{n-1} q^{-i}$ during n phases (if the first column height in phase n is 1), while it is possible to earn $q \sum_{i=0}^{n-1} q^{-i}$ in these phases.*

Suppose ALG acts as in Observation 3 for N consecutive phases, starting from some column i_0 in state A . We will ignore all previous phases in our calculations and only give bounds for the competitive ratio of ALG on these N phases. If we find some lower bound of $\mathcal{R}' < q$ on these phases, this also holds for the entire input by Claim 4. Let $\text{ALG}(j)$ be the total profit of ALG after j phases, ignoring the profit ALG_0 which was obtained before these phases started, and scaled such that the first column in phase j has height 1. Define α_j by the equality

$$\text{ALG}(j) = \sum_{i=1}^j x_i q^{i-j} = \alpha_j \sum_{i=0}^{j-1} q^{-i}. \quad (5)$$

Since $0 < x_i \leq 1$ for all $i \geq 1$, we have

$$0 < \alpha_j \leq 1 \text{ for all } j. \quad (6)$$

Since we scale in each step, and the column heights grow by a factor of q in each phase, we have

$$\text{ALG}(j+1) = \frac{\text{ALG}(j)}{q} + x_{j+1}. \quad (7)$$

We will use repeatedly that for $j \geq 1$,

$$\sum_{i=0}^{j-1} q^{-i} = \frac{1 - q^{-j}}{1 - q^{-1}} = \frac{q - q^{1-j}}{q - 1}. \quad (8)$$

Lemma 8 *Assume that ALG acts as in Observation 3 for N consecutive phases and has a competitive ratio less than $(1 - \varepsilon)\mathcal{R}$. Then for all $0 < \varepsilon < 0.2$ there exists an N' (independent of N) such that for all $N' \leq j \leq N$, we have:*

- (i) $x_j < (1 - \varepsilon)\alpha_{j-1}$, and
- (ii) $\alpha_j < (1 - \frac{\varepsilon}{4})\alpha_{j-1}$.

Since these two statements lead to a contradiction for N large enough (by (ii) we get $\alpha_j \rightarrow 0$ for $j \rightarrow \infty$ and hence by (i) also $x_j \rightarrow 0$, contradicting that $x_j > 1/2$ by Lemma 7), this proves the lower bound for this type of algorithms.

Proof (i) We consider the input where a final column of height $x'_j = (1 - \delta)x_j$ arrives (for some $\delta > 0$) after the last column of phase j , which has height q . This is state change II in Figure 7. Now, since $x_j \leq 1$ it is possible to earn $4x'_j = 4(1 - \delta)x_j$ on the last four columns of this modified input:

Input	q^{1-j}	q^{2-j}	q^{2-j}	q^{3-j}	...	q^{-1}	1	1	q	x'_j
State	A	B	A	B	...	A	B	A	D	-
Phase	1		2		...	$j-1$		j		
ALG	x_1	0	x_2	0	...	x_{j-1}	0	x_j	x_j	0
OPT	0	q^{2-j}	0	q^{3-j}	...	0	x'_j	x'_j	x'_j	x'_j

This is the only case in which the adversary deviates from its standard strategy (namely in phase $j-1$ and j). Again we use the best possible strategy for ALG (given that its assignment in phases $1, \dots, j$ is fixed). Letting $\delta \rightarrow 0$, the competitive ratio for this input is

$$\frac{4x_j + q \sum_{i=2}^{j-1} q^{-i}}{2x_j + \alpha_{j-1} \sum_{i=1}^{j-1} q^{-i}} = \frac{4(q-1)x_j + 1 - q^{2-j}}{2(q-1)x_j + \alpha_{j-1}(1 - q^{1-j})}.$$

(Note the indices of the summations.) For any $\varepsilon > 0$, there exists an N_1 such that $q^{2-j} < \varepsilon$ for all $j \geq N_1$. Then using (6) it can be verified that

$$\frac{4(q-1)(1-\varepsilon)\alpha_{j-1} + 1 - \varepsilon}{2(q-1)(1-\varepsilon)\alpha_{j-1} + \alpha_{j-1}(1-\frac{\varepsilon}{q})} > (1-\varepsilon)\mathcal{R}_1.$$

(We get equality for $\alpha_{j-1} = 1/(1-4.67\varepsilon)$, but by (6) this cannot happen. Note that $1-4.67\varepsilon > 0$ since $\varepsilon < 0.2$.) Since we can select δ arbitrarily small, we must have $x_j < (1-\varepsilon)\alpha_{j-1}$ for all $j \geq N_1$.

(ii) We first verify that

$$\frac{\alpha_j}{q} \cdot \frac{q}{q-1} + (1-\varepsilon)\alpha_j < \left(1 - \frac{\varepsilon}{2}\right) \alpha_j \cdot \frac{q}{q-1}$$

for $q = 2.144472$, for any $\alpha_j > 0$ and $\varepsilon > 0$ (the inequality reduces to $q/(2q-2) < 1$). Using (5), (7) and (i), this shows that

$$\text{ALG}(j+1) = \frac{\alpha_j}{q} \sum_{i=0}^{j-1} q^{-i} + x_{j+1} < \left(1 - \frac{\varepsilon}{2}\right) \alpha_j \cdot \frac{q}{q-1}$$

for $j \geq N_1 + 1$. By (5), for every $\varepsilon > 0$ there exists an N_2 such that $\text{ALG}(j+1) > \alpha_{j+1} \frac{q}{q-1} / (1 + \varepsilon/4)$ for all $j \geq N_2$. Hence for all $j \geq \max(N_1 + 1, N_2)$, we get

$$\alpha_{j+1} < \left(1 - \frac{\varepsilon}{2}\right) \alpha_j \left(1 + \frac{\varepsilon}{4}\right) < \left(1 - \frac{\varepsilon}{4}\right) \alpha_j.$$

We can now define $N' = \max(N_1 + 1, N_2)$. □

It remains to be shown that ALG cannot do better by acting differently than in Observation 3. Given Corollary 1, we only need to deal with phases of the form $A^k B$ for some $k \geq 1$. This is done by modifying the input as soon as such a phase occurs after there have been sufficiently many previous phases (of any length). As stated, we change q to $q' = 2.03644$ and p to $p' = 1/q'$ (at the point where this phase of length $k > 1$ ends) and continue the input as before.

We will show that ALG now has no choice but to allocate a nonzero value in each state A column, and moreover that this value is decreasing over time. On a 0 allocation, state change I follows, and given that $p' = 1/q'$, it is now best to allocate 1 in state C . Eventually we find a contradiction to Lemma 7.

The calculations can be checked using the following table for $k = 3$. We number the phases so that the first phase with the modified values of p and q is phase 1. The first column in phase 1 has height 1. Phase 0 was the phase of length $k > 1$, which happened after many previous phases where OPT earned almost $1/(q^3 - q) = 0.12957$ (at least) due to Claim 5. (To be precise, we wait until OPT has earned at least 0.129560 on previous phases; this happens already after at most 12 columns, independently of the number of phases.) Using Claim 4, we use $0.12956/q$ as an upper bound for $\text{ALG}(-1)$; to this we add $1/q$ for phase 0. Generally, we use $\text{ALG}(i-1) \leq \text{OPT}(i-1)/q$. In all calculations below, if $\text{OPT}(i-1)$ is larger than the lower bound that we use, and $\text{ALG}(i-1) = \text{OPT}(i-1)/q$, the overall competitive ratio only increases.

In the table, h_i is the height of the newly arriving column in phase i (so $h_0 = 1$). We verify that ALG can never allocate 0 in the column labeled Ratio after 0 allocation: in this case we have $\text{ALG}(i) = h_i + \text{ALG}(i-1)$, $\text{OPT}(i) = 4h_{i-1} + \text{OPT}(i-1) - h_{i-1}$. The allocation x_i is given by the formula $x_i = (\mathcal{R} \cdot \text{ALG}(i-1) - (\text{OPT}(i-1) - h_{i-1})) / (4 - 2\mathcal{R})$ so that an allocation above x_i immediately leads to a competitive ratio of \mathcal{R} .

Assuming an allocation below x_i , we find $\text{ALG}(i) \leq \text{ALG}(i-1) + x_i h_{i-1}$.

Phase	$\text{ALG}(i-1)$	$\text{OPT}(i-1) - h_{i-1}$	h_i	Ratio after 0 allocation			x_i	$x_i h_{i-1}$
1	0.5267	0.3470	2.0364	4.3470 /	2.5632	= 1.696	0.8984	0.8984
2	1.4251	1.3470	4.1471	9.4928 /	5.5722	= 1.7036	0.8640	1.7594
3	3.1845	3.3835	8.4453	19.972 /	11.630	= 1.7173	0.7999	3.3173
4	6.5018	7.5306	17.198	41.312 /	23.70	= 1.7431	0.6808	5.7493
–	18.000	33.174						

As an example, we show the two ways in which the input may end on the first line of the above table. Let $x'_1 = (1 - \delta)x_1$.

Input	$\dots q^{-3}$	q^{-3}	q^{-2}	q^{-1}	1	1	q'	1
State	$\dots B$	A	A	A	B	A	C	–
Phase	$\dots, -2, -1$	0				1		
ALG	0.0604	0	0	q^{-1}	0	0	q'	0
OPT	0.12956	0	q^{-2}	0	1	1	1	1
Input	$\dots q^{-3}$	q^{-3}	q^{-2}	q^{-1}	1	1	q'	x'_1
State	$\dots B$	A	A	A	B	A	D	–
Phase	$\dots, -2, -1$	0				1		
ALG	0.0604	0	0	q^{-1}	0	\mathbf{x}_1	\mathbf{x}_1	0
OPT	0.12956	0	q^{-2}	0	x'_1	x'_1	x'_1	x'_1

For $k = 2$, we find $\text{OPT}(0) - h_0 = \text{OPT}(-1) + 1/q^2 = 0.4953$ (for a long enough input) and $\text{ALG}(0) \leq 1/q + \text{OPT}(-1)/q = 0.5959$. Hence, before the first column of height 1, OPT earns more relative to ALG than in the case $k = 3$. Since the profit ratio on the remaining part of the input remains unchanged and is always more than \mathcal{R} (either $4/q'$ or $2(1 - \delta)$ in the first line), we get the same results as before.

Similarly, for $k > 3$ it can be verified that OPT always earns more relative to ALG on early columns than in the case $k = 3$, leading to the same conclusion as above. (Increasing k by 2 always increases the relative profit of OPT.) Hence, after at most 12 columns of the input, ALG can only use phases of the form AB , which by Lemma 8 eventually leads to a competitive ratio arbitrarily close to $\mathcal{R} = 1.69595$.

5 Conclusions

We have narrowed the gap for this problem to 0.056. We believe that both our lower bound and upper bound could potentially be improved, but we conjecture that the lower bound is closer to the true competitive ratio of the problem. However, it is not easy to see how to narrow the gap further. There are four cases where the analysis for our algorithm is tight; additionally, there are various cases where the analysis is nearly tight. An improved algorithm would have to achieve a better ratio in all of the very different tight cases without losing too much in other cases.

Acknowledgment The author would like to thank Leah Epstein and Asaf Levin for interesting discussions, and Marek Chrobak for pointing out some problems in the proof of the lower bound in the conference version.

References

- [1] Amrinder Arora and Hyeong-Ah Choi. Channel aware scheduling in wireless networks. Technical Report 002, George Washington University, 2006.
- [2] Amrinder Arora, Fanchun Jin, and Hyeong-Ah Choi. Scheduling resource allocation with timeslot penalty for changeover. *Theor. Comput. Sci.*, 369(1-3):323–337, 2006.
- [3] Sem C. Borst. User-level performance of channel-aware scheduling algorithms in wireless data networks. In *INFOCOM*, 2003.
- [4] Leah Epstein and Asaf Levin. Private communication, 2009.
- [5] W. Stallings. *Wireless Communication & Networks*. Prentice Hall, 2001.
- [6] Vagelis Tsibonis, Leonidas Georgiadis, and Leandros Tassiulas. Exploiting wireless channel state information for throughput maximization. In *INFOCOM*, 2003.
- [7] Haitao Wang, Amitabh Chaudhary, and Danny Z. Chen. Online rectangle filling. In Christos Kaklamanis and Martin Skutella, editors, *WAOA*, volume 4927 of *Lecture Notes in Computer Science*, pages 274–287. Springer, 2007.
- [8] Haitao Wang, Amitabh Chaudhary, and Danny Z. Chen. New algorithms for online rectangle filling with k-lookahead. In Xiaodong Hu and Jie Wang, editors, *COCOON*, volume 5092 of *Lecture Notes in Computer Science*, pages 385–394. Springer, 2008.